

World Journal of *Medical Genetics*

World J Med Genet 2016 February 27; 6(1): 1-16





REVIEW

- 1 Gene regulatory networks in atrial fibrillation

Franco D, Lozano-Velasco E, Aranega A

ABOUT COVER

Editorial Board Member of *World Journal of Medical Genetics*, George Sai-Wah Tsao, PhD, Professor, Department of Anatomy, LKS Faculty of Medicine, University of Hong Kong, Hong Kong, China

AIM AND SCOPE

World Journal of Medical Genetics (*World J Med Genet*, *WJMG*, online ISSN 2220-3184, DOI: 10.5496) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJMG covers topics concerning genes and the pathology of human disease, molecular analysis of simple and complex genetic traits, cancer genetics, epigenetics, gene therapy, developmental genetics, regulation of gene expression, strategies and technologies for extracting function from genomic data, pharmacological genomics, genome evolution. The current columns of *WJMG* include editorial, frontier, diagnostic advances, therapeutics advances, field of vision, mini-reviews, review, topic highlight, medical ethics, original articles, case report, clinical case conference (Clinicopathological conference), and autobiography.

We encourage authors to submit their manuscripts to *WJMG*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

World Journal of Medical Genetics is currently no indexing/abstracting.

FLYLEAF

I-IV Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Ya-Jing Lu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Shui Qiu*
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL
World Journal of Medical Genetics

ISSN
ISSN 2220-3184 (online)

LAUNCH DATE
December 27, 2011

FREQUENCY
Quarterly

EDITOR-IN-CHIEF
Hans van Bokhoven, Professor, PhD, Department of Human Genetics and Cognitive Neurosciences, Radboud university Nijmegen Medical centre, PO Box 9101, 6500 HB Nijmegen, The Netherlands

EDITORIAL OFFICE
Jin-Lei Wang, Director
Xiu-Xia Song, Vice Director
World Journal of Medical Genetics

Room 903, Building D, Ocean International Center, No. 62 Dongsihuan Zhonglu, Chaoyang District, Beijing 100025, China
Telephone: +86-10-59080039
Fax: +86-10-85381893
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.wjgnet.com/esp/helpdesk.aspx>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
8226 Regency Drive,
Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.wjgnet.com/esp/helpdesk.aspx>
<http://www.wjgnet.com>

PUBLICATION DATE
February 27, 2016

COPYRIGHT

© 2016 Baishideng Publishing Group Inc. Articles published by this Open Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS

Full instructions are available online at http://www.wjgnet.com/bpg/g_info_20160116143427.htm

ONLINE SUBMISSION

<http://www.wjgnet.com/esp/>

Gene regulatory networks in atrial fibrillation

Diego Franco, Estefanía Lozano-Velasco, Amelia Aranega

Diego Franco, Estefanía Lozano-Velasco, Amelia Aranega, Cardiovascular Development Group, Department of Experimental Biology B3-362, Faculty of Experimental Sciences, University of Jaén, 23071 Jaén, Spain

Author contributions: Franco D wrote the manuscript; Lozano-Velasco E and Aranega E critically read the manuscript.

Supported by A CNIC translational grant (CNIC2006/08) to Franco D; and the University of Jaén on translational biomedicine to Franco D (UJA2013/01).

Conflict-of-interest statement: The authors declare that they have not conflict of interest.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Correspondence to: Dr. Diego Franco, Professor, Cardiovascular Development Group, Department of Experimental Biology B3-362, Faculty of Experimental Sciences, University of Jaén, CU Las Lagunillas s/n, 23071 Jaén, Spain. dfranco@ujaen.es
Telephone: +34-953-212763
Fax: +34-953-211875

Received: September 29, 2015
Peer-review started: October 21, 2015
First decision: November 27, 2015
Revised: January 14, 2016
Accepted: February 16, 2016
Article in press: February 18, 2016
Published online: February 27, 2016

Abstract

Atrial fibrillation (AF) is the most frequent arrhythmogenic syndrome in humans. With an estimate incidence of 1%-2% in the general population, AF raises up to almost

10%-12% in 80+ years. Thus, AF represents nowadays a highly prevalent medical problem generating a large economic burden. At the electrophysiological level, distinct mechanisms have been elucidated. Yet, despite its prevalence, the genetic and molecular culprits of this pandemic cardiac electrophysiological abnormality have remained largely obscure. Molecular genetics of AF familial cases have demonstrated that single nucleotide mutations in distinct genes encoding for ion channels underlie the onset of AF, albeit such alterations only explain a minor subset of patients with AF. In recent years, analyses by means of genome-wide association studies have unraveled a more complex picture of the etiology of AF, pointing out to distinct cardiac-enriched transcription factors, as well as to other regulatory genes. Furthermore a new layer of regulatory mechanisms have emerged, *i.e.*, post-transcriptional regulation mediated by non-coding RNA, which have been demonstrated to exert pivotal roles in cardiac electrophysiology. In this manuscript, we aim to provide a comprehensive review of the genetic regulatory networks that if impaired exert electrophysiological abnormalities that contribute to the onset, and subsequently, on self-perpetuation of AF.

Key words: Atrial fibrillation; Genetics; MicroRNAs; Genome-wide association studies; PITX2

© The Author(s) 2016. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Atrial fibrillation (AF) is the most prevalent arrhythmogenic defect in the human population. Genetic factors such as mutations in distinct ion channel encoding genes have been described, yet representing less than 10% of all AF cases. Genome wide association studies have widened the genetic culprits contributing to AF. We provide herein a state-of-the art review on the genetic components underlying AF. Experimental evidences demonstrated that PITX2 plays a pivotal role regulating cellular, molecular and electrophysiological characteristics of the developing and adult heart that, if impaired, predispose to AF, leading to complex regulatory networks with transcriptional and post-transcriptional (microRNA)

regulatory mechanisms.

Franco D, Lozano-Velasco E, Aranega A. Gene regulatory networks in atrial fibrillation. *World J Med Genet* 2016; 6(1): 1-16 Available from: URL: <http://www.wjgnet.com/2220-3184/full/v6/i1/1.htm> DOI: <http://dx.doi.org/10.5496/wjmg.v6.i1.1>

BACKGROUND

Atrial fibrillation (AF) is the most frequent arrhythmogenic syndrome in humans. With an estimate incidence of 1%-2% in the general population, AF raises up to almost 10%-12% in 80+ years^[1-5]. Cardiovascular risk factors such diabetes, obesity, hypertension and hyperthyroidism promote AF^[6-9]. In addition, the occurrence of AF can be also triggered by preceding cardiovascular diseases such as hypertrophic cardiomyopathy and valvular heart diseases^[10,11] as well as it can be boosted by cardiac surgery, obstructive sleep apnea and inflammatory processes^[12-15]. Importantly, besides the global risk factors for AF, it is well-established that the debut of an AF episode triggers subsequent and more severe AF episodes, leading to electrical and structural remodeling of the diseased heart, a condition quoted as "AF begets AF"^[16,17]. Thus AF *per se* is a progressive disease. Electrical remodeling involves progressive changes in the cardiac electrical properties, possibly triggered by oxidative stress imbalance, leading to EADs, DADs and/or changes in the ADP configuration^[18,19], culminating thus in rotor formation^[20] (Figure 1). Structural remodeling involves atrial dilation, fibrosis and/or inflammation^[21,22] which then indirectly promotes the generation of rotors and thus electrical re-entry circuitries^[20] (Figure 1). Moreover, suffering from AF predisposes to additional health problems, ranging from bleeding and stroke^[23-25], cognitive decline and dementia^[26,27], ventricular dysfunction^[28,29] and even sudden death^[30] (Figure 1). Thus, these data support the notion that AF is an epidemic disease with large socio-economic burden worldwide.

While large epidemiological data support the previously mentioned predictive risk factors for AF, there are also unquestionable evidences that in a subset of AF patients no concurrent previous risk factors are diagnosed and thus AF occurs in an isolated manner, a condition dubbed lone AF^[31-34]. These observations, in addition to seminal observations supporting that AF can also be familial^[35], strengthen the notion of a genetic component on the onset of AF. In this context, two waves of genetic information have been gained over the last decades. Seminal studies on familial AF cases followed by genetic linkage analyses, and subsequently by genetic screening of candidate genes identified a large number of point mutations in distinct genes encoding for proteins involved in cardiac electrophysiology^[36-38]. These data, along with meticulous electrophysiological mapping analyses and *in silico* data modeling, provided an important cornerstone to understand AF pathophysiology^[39-43].

However, genetic identification of candidate AF genes can only explained around 10%-15% of all AF patients. Thus, new approaches to untangle the genetic bases of AF were envisioned. Genome-wide association analyses (GWAS) lightened the discovery of new genes associated to AF. Seminal worked by Gudbjartsson *et al.*^[44] firstly identify common risk variants highly associated to the onset of lone AF in distinct large European and Chinese patient cohorts. Subsequent GWAS studies and meta-GWAS have further identified new candidate genes for AF pathophysiology^[45-49], yet the functional link between these variants and the AF pathophysiology is a rather unexplored. A tortuous route to dissect their functional roles is envisioned, given that most of these risk variants are located rather apart from gene coding sequences. Thus, a first proposal based on these findings is that risk variants might affect gene regulatory networks within the vicinity and thus underscore AF onset. In addition, a novel layer of complexity in gene regulatory networks has emerged with the discovery of non-coding RNAs that can mediate post-transcriptional regulation. In this review, we aim to elaborate on the current state-of-the-art of the gene regulatory networks involved in AF pathophysiology.

CONTRIBUTION OF THE IMPAIRED CARDIAC ACTION POTENTIAL TO THE AF PATHOPHYSIOLOGY

The cardiac action potential is driven by inward and outward flow of distinct ion currents along the cardiomyocyte membrane. In atrial and ventricular cardiomyocytes, the configuration of the action potential is initiated by the upstroke depolarization of the sodium current, followed by a repolarization mediated by distinct potassium currents until the resting membrane potential is restored. A conduction-contraction coupling is modulated within each action potential by a complex regulatory network of calcium handling^[50]. Interestingly, nodal conductive cells display a distinct upstroke configuration which is modulated by cation selective currents, yet repolarization is rather similar to working cardiomyocytes. A large number of mutations in the *SCN5A* gene, coding for the pore-forming subunits of the I_{Na} current, have been associated to AF, alone^[51-56] or in combination to other concurrent cardiac pathologies^[57-62]. Similarly, mutations in *SCN1B*, *SCN2B*, *SCN3B* and *SCN4B* ancillary subunits, respectively, have also been linked to AF^[63-67]. At present, the functional interpretation of these findings suggest that impaired I_{Na} current can promote AF onset, yet electrophysiological evidences are only available for a subset of these mutations. Importantly, mutations in *SCN5A* are also linked to other cardiac arrhythmias such as long QT and Brugada^[68-70] demonstrating a pivotal role for *SCN5A* on cardiac electrophysiology and illustrating that distinct phenotype might be acquired depending on subtles differences on the mutation location as well as on plausible modulatory accompanying proteins. In addition, mutations in the *HCN4* cation channel, responsible for the cardiac




Figure 1 Schematic representation of the clinically relevant risk factors influencing the onset of atrial fibrillation. Atrial fibrillation (AF) leads thereafter to distinct cellular and molecular remodeling events, including structural and electrical remodeling, in part as caused by inflammatory process. In addition, progression of AF is highly associated with the onset of additional cardiac pathophysiology.

action potential upstroke of conductive cells, have also been reported in AF^[71].

Mutations in genes encoding for proteins controlling the repolarization phase of the cardiac action potential have also been associated to AF. In particular, *KCNQ1*^[36,72-79], *KCNAS*^[80] and *KCNE2*^[38,81] mutations have been identified. In this setting, point mutations seem to shorten the repolarization phase providing an electrophysiological substrate for AF onset. Similarly, alteration of the resting membrane potential have been related to trigger AF onset and a large number of mutations in the gene coding for *KCNJ2* have been reported^[37,38,82,83]. Similarly as for the sodium channels, mutation in potassium channels have been associated to AF and concomitantly with other arrhythmogenic syndromes such as long^[72] and short^[84] QT, respectively.

Multiple lines of evidences have demonstrated the essential role of impaired calcium homeostasis as a triggering factor of AF^[2,85-90]. In this context, mutations in *RYR2* are associated to AF^[91] and in conjunction with other cardiac pathophysiological conditions such as catecholaminergic polymorphic ventricular tachycardia^[92,93]. Curiously, *RYR2* mutations can also affect sodium channel expression^[94]. Importantly, murine model of *RYR2* mutation leads to AF, reinforcing the genetic evidences of impaired calcium homeostasis as a trigger of AF^[95].

In addition, mutations in gap junction proteins such as *GJA1* (connexin43)^[96,97] and *GJA5* (connexin40)^[98-104], which are critical for the coordinated transmission of the electrical impulse among cardiomyocytes, have also been identified in association to AF. Seminal work described that mutation in connexin proteins were occurring in somatic cells (*i.e.*, cardiomyocytes)^[98], whereas more recently germline mutations have also been identified^[98-104] and the concept of somatic cell mutation challenged^[105]. In this setting, electrophysiological analyses of connexin

mutations in the AF context demonstrated that cardiomyocyte electrical transmission is impaired^[106-111] and thus can result in cardiac re-entry circuitries and then on AF.

Furthermore, mutations in several other genes, besides those directly contributing to the electrophysiological properties of the heart, have also been identified in the AF setting. In particular, mutations in the transcription factors *TBX5*^[112], *NKX2.5*^[113-117] and *NKX2.6*^[118] as well as in the nuclear pore component *NUP155*^[119] and the atrial natriuretic factor (*ANP*)^[120]. A summary of gene mutations associated to AF is provided on Figure 2. It is important to highlight in this context that mutation in *NUP155* was found in patients with AF and early sudden death^[118], while *TBX5* and *NKX2.5* are developmental transcription factors that play determinant roles during cardiogenesis and mutations in these genes have also been reported in distinct congenital heart diseases as well as GWAS candidate genes in other cardiac electrophysiological defects^[121-128].

NOVEL GENES LINKED TO AF

As previously said, classical genetic approaches provided an entry site to discover discrete genes involved in AF pathophysiology, but felt short to explain most of the diagnosed AF cases. Genome-wide association studies (GWAS) introduced a revolutionary genetic approach to understand AF, and cardiac arrhythmias at large^[129,130]. Seminal work by Gudbjartsson *et al.*^[44] identified risk variant at 4q25 highly associated to lone AF. Soon thereafter these observations were corroborated in distinct studies worldwide^[131-136], yet with some controversial findings^[137,138]. Surprisingly, 4q25 risk variants are located in a gene desert, being the closest annotated gene a homeobox transcription factor, *PITX2*, around 150 kb downstream. Soon thereafter, experimental observations provided




Figure 2 Schematic representation genes associated to atrial fibrillation. On the one hand, gene mutations associated to atrial fibrillation (AF) are illustrated. On the other hand, genes associated to AF by means of GWAS are also illustrated. In the latter, three subset of genes are provided: (1) those directly linked to AF GWAS analyses; (2) those related to AF, but identified in arrhythmogenic syndromes GWAS analyses; and (3) risk variants association studies in small AF cohorts by a candidate approach strategy. Genes highlighted in bold represent those genes identified in AF by multiple approaches. GWAS: Genome-wide association analyses.

evidences that impaired *Pitx2* function in animal models triggered increase atrial arrhythmias susceptibility^[139,140]. However, how risk variants can influence *PITX2* expression and/or function have only been partially revealed^[141], since regulatory elements containing the risk variants can molecularly interact with *PITX2* but also *ENPEP* promoter elements in a tissue- and developmental-specific manner^[141]. Importantly, controversial findings in humans have been reported. Chinchilla *et al.*^[136] firstly described decreased *PITX2* expression in AF patients, yet these claims were challenged since no correlation between risk variants and *PITX2C* expression was observed^[142]. However, more recently a correlation with *PITX2A* expression has been reported, limiting their functional significance since data were obtained from right atrial biopsies^[143]. Thus, further investigations are required to clarify these controversial findings.

Besides *PITX2*, additional GWAS have enlightened the putative role of other novel genes in AF pathophysiology, such as *KCNN3*^[47], *ZFH3*^[45,48], *IL6R*^[49] and *MLY4*^[144]. In most cases, these findings have been corroborated in independent studies with distinct ethnicity cohorts as for *ZFH3*^[145,146], and *IL6R*^[147], but to date no additional reports are available for *KCNN3* and *MYL4*.

While arguments for the plausible involvement in AF have been provided in all cases, experimental evidences remain largely missing. *KCNN3* loss-of-function mutant mice display no overt cardiac electrophysiological defects, yet overexpression leads to sudden death^[148]. Similarly *ZFH3* function role in the heart is scarce, yet recent findings suggests that atrial arrhythmias^[149] and pacing^[150] leads to significant impaired *ZFH3* expression. Importantly, risk variants associated to *KCNN3*, *ZFH3* and *IL6R* lie within intronic regions, respectively, and thus have no predictive functional consequences. Therefore, as in the cases of 4q25 risk variants (*PITX2*), a functional role on plausible regulatory elements is hypothesized,

but functional evidences are yet missing.

In addition to these GWAS studies, at least two meta-GWAS studies have digged for additional risk variants associated to AF^[46,151]. In this context, six new loci have been uncovered using such approach; *CAV1*, *HCN4*, *SYNE2*, *SYNPOL2*, *PRRX1* and *WTN8A*. Replication analyses of the risk variants have been confirmed for *SYNE2*^[152] and *PRRX1*^[149], are controversial for *CAV1*^[53,153,154] or have not been reported to date for *HCN4* and *SYNPOL2*. Importantly, a 12-SNPs risk score have been developed that can individually identify the risk of AF and stroke^[155].

While this approach broadens the plausible genetic bases of AF, it is also true that their contribution to explain AF pathophysiology would be directed linked to the statistical significance associated with those findings. It is important to say in this context that risk variants associated to *PITX2* display highest significance values, *KCNN3*, *ZFH3* and *IL6R* are moderately significant and those revealed by meta-GWAS display the lowest, yet statistical significant, values. In a simplistic way, this could be transferred as *PITX2* being contributing to larger spectrum of AF pathophysiology and thus being in the upper level of a hierarchical signaling pathway while *CAV1*, *HCN4*, *SYNE2*, *SYNPOL2*, *PRRX1* and *WTN8A* will be rather discretely contributing to AF pathophysiology in a rather discrete, small subset of AF patients, and therefore being at the bottom of such hierarchy. We have recently provided experimental evidences supporting this notion, as detailed below.

GWAS studies have also been reported in other electrophysiological parameters such as PR interval^[129,156-158], heart rate^[157], QRS interval^[157] as well as physiopathological conditions such as Brugada syndrome^[159] and long QT^[160-162]. Several of these cardiac arrhythmogenic conditions are sometimes directly associated to AF, such as for example impaired PR interval or Brugada syndrome. Importantly,

such GWAS studies have identified multiple genes, some of which are shared with AF, such as *CAV1* (PR interval^[129,156-158]) while other have been previously involved in AF pathophysiology, such as *KCNQ1*, *KCNH2*, *SCN5A*, *KCNJ2* (long QT GWAS analyses^[160-162]). In addition to GWAS studies, risk variants have been associated to AF using more discrete/small AF patient cohorts. Among them it is important to highlight risk variants at the following loci; *KCNQ1*^[163-165], *KCNH2*^[163], *KCNE1*^[163], *KCNJ5*^[166], *GJA5*^[167] and *TBX5*^[168]. In addition to these GWAS studies, several independent AF cohort studies have reported risk variants associated to AF in other loci, such as *TNNI3*^[169] and *SCN10A*^[170,171], yet the functional relevance of these findings remains unsolved. These data suggest that complex and intricate gene networks are likely to be operative between these regulatory genes resulting in AF (Figure 3) as well as other electrophysiological associated disorders.

A NOVEL AND EMERGING LAYER OF GENE REGULATORY CONTROL

Gene regulatory networks have been classically associated as hierarchical interactions between master genes, such as transcription factors, and structural genes, such as actin and myosin genes forming the sarcomere or genes encoding ion channels configuring therefore the cardiac action potential. In this setting, transcriptional regulation has been considered the master regulatory point, deciding whether to go or not to go, whereas translation and post-translational modifications are considered as minor regulatory nodes. Over the last decade, we have witnessed the discovery that large part of the so called "rubbish DNA" in fact codes for a large variety of non-coding RNAs species with highly relevant functional properties. In broad sense, non-coding RNAs can be sub-classified into long non-coding (lncRNAs) and small non-coding RNAs, exerting a distinct and variable functional roles^[172]. Among small non-coding RNAs, microRNAs have emerged as the larger group, playing a pivotal role on post-transcriptional regulation of mRNA transcript expression^[173]. MicroRNAs are 22-24 nt non-coding RNAs which can block translation or trigger transcript degradation by binding to complementary seed sequence in the 3'untranslated region (3'UTR) of nascent mRNA transcripts^[173]. At present the hallmark of microRNA transcriptomics has been uncovered in the developing and adult heart in both normal and pathological conditions^[174,175]. In the context of AF the microRNA fingerprint has been profiled in different settings^[176-178], providing highly valuable biomarkers^[179-183] as well as opening new avenues to dissect the functional role of these differentially expressed microRNAs in AF pathology. In addition distinct sets of microRNAs have also been reported to selectively modulate expression of distinct gene encoding for ion channels, opening thus new pathways to dissect their genetic contribution to AF.

AN INTEGRATIVE GENE REGULATORY NETWORK LINKED TO AF

As stated above, evidence on the functional role of multiple genes have been provided by the identification and experimental testing of AF patient gene mutations. In this setting, a causative link can, in most cases, be established. However, GWAS analyses provide circumstantial evidences whereas functional linkage between the risk variants and coding genes, whether protective or deleterious, is not directly established. To date, an example of successful linkage have been reported^[184] in the cardiovascular context, however in most cases, this is not yet established. Therefore, dissecting the functional role of GWAS candidate genes represents the first step for the analyses of their causative link.

Large set of evidences suggest that multiple genetic pathways can contribute to the onset of AF. Over the last years we have carefully dissected the role of the homeobox transcription factor *PITX2* in this context and we have made efforts to dissect the regulatory interaction of *PITX2* with those genes previously reported to play a role in AF pathophysiology. Several studies have demonstrated that *PITX2* can influence expression of distinct ion channels in the developing and adult hearts. Wang *et al.*^[138] firstly demonstrate that *PITX2* haploinsufficiency is sufficient to increase AF susceptibility. These authors demonstrated that *PITX2* haploinsufficiency led to developmental impaired expression of key transcription factors (*SHOX2*, *TBX3*) directing sinoatrial morphogenesis and thus impacting on the normal onset of the cardiac pacemaker structures. However, in this context, no basal AF occurs and no ECG defects were observed. In this line of thinking soon thereafter Kirchoff *et al.*^[139] demonstrate similar findings and furthermore reported that expression of multiple ion channels were impaired as well as cell-cell interacting proteins (Figure 3). However, none of these genes were reported to play a pivotal role on cardiac action potential and thus, molecular and electrophysiological links were still on its infancy. Using a distinct experimental approach, *i.e.*, an atrial-specific *PITX2* conditional transgenic mouse line, Chinchilla *et al.*^[135] revealed that ECG defects were observed at rest (no triggering needed), electrophysiological measurements were impaired, *i.e.*, increased resting membrane potential and prolonged action potential duration, and furthermore using a candidate approach analyses, distinct ion channels were also impaired, particularly *SCN5A* (I_{Na}), *KCNJ2* (I_{K1}) and *KCNH2* (I_{Kr}). Thus, these data demonstrate for the first time that functional impairment of *PITX2* leads to molecular defects which provoke electrophysiological alterations and thus increase rate of atrial arrhythmias. In line with these findings, Tao *et al.*^[185] using a *PITX2* deletion conditional approach in the adult heart also demonstrated that basal ECG defects were observed, similar to those already reported by Chinchilla *et al.*^[135], and moreover they




Figure 3 Schematic representation of the *Pitx2* downstream pathways involved in pro-arrhythmic events leading to atrial fibrillation. Multiple lines of evidences demonstrate that *PITX2* exerts a pivotal role regulating expression of distinct ion channels, cell-cell coupling, blood pressure controllers and beta-adrenergic stimulation. Most of these pathways are modulated by microRNAs which are under control of *PITX2* action. Red denotes down-regulation and green up-regulation.

demonstrated that key determinants of the cardiac action potential were impaired, such as *KCNQ1* (I_{Ks}), *KCNH2* (I_{Kr}), *CACNA1C* (I_{Ca}), among others. Overall these data demonstrated that *Pitx2* insufficiency can both promote and trigger molecular and electrophysiological alterations, namely affecting the depolarization, repolarization and resting membrane potential characteristics of the cardiac action potential. They also illustrate that alteration of a single transcription factor can influence the functional role of multiple ion channels previously reported to be associated to AF. Therefore it can be inferred that *PITX2* is upstream of this hierarchical signaling pathway and its impairment would explain a large fraction of AF patients.

In addition to *PITX2*, other GWAS studies have provided evidence on the plausible role of distinct gene in AF pathophysiology, yet as previously said, their functional implication has been poorly documented. We investigated whether these genes would be under control of *PITX2*. Using a series of gain and loss-of-function approaches, Lozano-Velasco *et al.*^[149] demonstrated that *PITX2* can modulate expression of *ZFH3*, *KCNN3* and *IL6R*. The

role of *ZFH3* in atrial arrhythmias is basically unexplored. However, given its transcriptional capacity it might be plausible that it can modulate multiple genes previously related to AF. The role of *KCNN3* in atrial arrhythmias is under debate, its plausible contribution to the configuration of the cardiac action potential, if any, seems to be residual. On the other hand, the involvement of *IL6R* could be related to inflammatory processes that, if impaired, are associated to AF as previously mentioned. We have noticed that *IL6R* expression in the atrial chambers is impaired in *PITX2* deficient mice yet plasma levels of soluble *IL6* and *TFN α* ligands are not altered. While these data support a tissue-specific involvement of *IL6R*, its links to AF remain to be elucidated.

We also investigated if *PITX2* can influence the role of those genes related to AF by meta-GWAS analyses. Among them, previous evidences support that *PITX2* can modulate expression of *HCN4*^[138]. More recently, Lozano-Velasco *et al.*^[151] demonstrated that also *CAV1*, *SYNE2* and *WNT8A* are modulated by *PITX2*. It is important nonetheless to highlight that risk variants associated

to *CAV1* and *SYNE2* have also been reported in other cardiac electrophysiological disorders yet their functional role remains poorly characterized. Importantly, WNT signaling has been largely documented to play pivotal role during cardiac development and homeostasis^[186,187], however WNT8 has not been reported to play fundamental roles therein^[188].

Given the potential signaling properties of WNT8A and those transcriptional capacity exerted by *ZFH3* we explored in any of those pathways might be involved on the regulation of distinct ion channels previously reported to be modulated by *PITX2* and related to AF. Gain- and loss-of-function approaches demonstrated that *WNT8A* but not *ZFH3* can modulate expression of calcium handling proteins^[151]. Importantly, impaired expression of *WNT8A* is observed in *PITX2* insufficiency models with basal ECG alterations but not in those with normal ECG recordings^[151], supporting the notion that WNT signaling plays a fundamental role conferring susceptibility vs triggering capacity to the atrial chambers during AF onset.

In addition to gene regulatory mechanisms driven by *PITX2*, several lines of evidence support that *PITX2*-regulated microRNAs also contribute to AF pathophysiology. Chinchilla *et al*^[135] already demonstrated that impaired miR-1 expression in *PITX2* deficient mice underlie abnormal resting membrane potential configuration (by modulating *KCNJ2* and *KCNJ12* post-transcriptional regulation). Huang *et al*^[153] has confirmed that *PITX2* inhibits miR-1 expression, which negatively regulates *ZFH3*. Wang *et al*^[189] demonstrated that two distinct genomic microRNA clusters, namely *miR-17-92* and *miR-106b-25*, as well as *miR-335* and *miR-423*, are under control of *PITX2*. Genetic deletion of *miR-17-92* and *miR-106b-25* microRNA clusters, respectively display normal baseline electrophysiological parameters but were susceptible to pacing-induced AF. Cardiac-specific deletion (Nkx2.5Cre) of *miR-17-92* cluster resulted in prolonged PR interval. Compound double mutant mice using conditional Nkx2.5Cre mediated deletion resulted in sinus node dysfunction, a condition reminiscent of sick sinus syndrome in humans. Molecular analyses revealed that *SHOX2* and *TBX3* expression were impaired. Furthermore, RYR2-mediated calcium-leak was reported in *miR-106b-25* deficient mice^[190] since several members of this cluster, namely miR-106b, miR-93 and miR-25 directly target the 3' UTR of Ryr2. More recently, Lozano-Velasco *et al*^[149] demonstrate that a large number of microRNAs, previously reported to be associated to AF in patients, are also modulated by *PITX2*. In particular, *miR-1*, *miR-26b*, *miR-29a*, *miR-30e*, *miR-106b*, *miR-133* and *miR-200* were up-regulated in absence of *PITX2* whereas *miR-21*, *miR-106a*, *miR-203*, *miR-208a* and *miR-208b* were down-regulated. Several of these microRNAs, such as *miR-1*, *miR-133*, *miR-21*, *miR-106b* and *miR-26* have been previously reported to regulate calcium (*CACNA1C*^[191]; *RYR2*^[190]), sodium (*SCN5A*^[192]), potassium (*KCNJ2*^[193], *KCNE1*^[194], *KCNB2*^[194]), cation (*HCN4*^[195]) channel subunits, respectively. Overall, these

data demonstrate a highly complex gene regulatory networks leading to AF as summarized in Figure 3.

CONCLUSION

In this review we have highlighted the complexity of risk factors influencing the onset of AF, both clinically and genetically. Several lines of evidences demonstrated that *PITX2* exerts a pivotal role on the genetic determinants of AF^[136,139,140,149,185]. While it has been robustly demonstrated that *PITX2* insufficiency predisposes to atrial arrhythmia genesis in experimental mouse models^[136,139,140,149,185], discrepancies remain as whether *PITX2* is impaired in AF risk variants human patient carriers^[136,142,143]. Importantly some discrepancies also exist as whether predisposes or triggers AF^[136,139]. Our recent findings suggest the notion that *PITX2* insufficiency in the embryo predisposes whereas *PITX2* insufficiency in the adult atrial chamber triggers AF. In this context, we have provided evidences that such switching mechanisms seems to be regulated by WNT signaling and the downstream activation/repression of key microRNAs^[149,165], yet future experiments are needed to provide additional evidences on this front. In the last decade, our understanding of the downstream pathways controlled by *PITX2* has advanced at a quick pace. Several studies have demonstrated that *PITX2* controls signaling pathways regulation sinus node formation^[189,196], including therein the pivotal role of a microRNA cluster^[189,190]. These data shed light thus on the concurrence of AF with other cardiac physiopathological conditions such as sick sinus syndrome^[190]. Similarly we and others have demonstrated the key role exerted by *PITX2* regulating microRNAs impacting on calcium handling and thus on profibrillatory risk factors^[136,153,189]. It remains nonetheless to be established if *PITX2* also leads to impaired regulation of inflammatory processes and/or redox signaling which might impact on structural remodeling of the atrial chambers. To date, no evidences of atrial fibrosis have been reported in any of the *PITX2* deficient mouse models^[136,139,140,149,185], while some incipient evidences have been reported on the inflammatory link^[149] but not on redox signaling. Thus, future lines of research should be envisioned to clarify this point.

In addition, whereas our understanding of the *PITX2* downstream signaling pathways in the context of AF have progressively increased, scarce insights are currently available on the impact of AF clinically related risk factors on *PITX2*. A link between 4q25 risk variant carriers and increase left atrial volume has been recently reported^[197] in AF patients, however it remains to be established this is modulated by *PITX2*, as suggested by previous evidence in *PITX2* deficient mice in which atrial volume is already increased at fetal developmental stages^[136]. A recent study reported that aging and hypertension, two well-established risk factors for AF, severely decreased *PITX2* expression in a rat experimental model when both risk factors were combined^[198]. We recently reported that *Pitx2* is severely impaired in dilated cardiomyopathy patients as well as in an experimental heart failure pig model, shedding light

into a possible connection between ventricular dysfunction and AF^[199,200]. Yet, it remains to be established if other clinically relevant AF risk factors, such as diabetes, obesity, hyperthyroidism, valvular heart disease and/or obstructive sleep apnea are also impairing *PITX2* function and thus predisposing to AF. Overall these data demonstrate the pivotal role of *PITX2* regulating multiple aspect that if impaired are pro-arrhythmic and they also open new pathways to explore therapeutical approaches that could eventually lead to minimize the burden of AF in the human population.

REFERENCES

- Hakim FA**, Shen WK. Atrial fibrillation in the elderly: a review. *Future Cardiol* 2014; **10**: 745-758 [PMID: 25495816 DOI: 10.2217/fca.14.32]
- Nattel S**, Dobrev D. The multidimensional role of calcium in atrial fibrillation pathophysiology: mechanistic insights and therapeutic opportunities. *Eur Heart J* 2012; **33**: 1870-1877 [PMID: 22507975 DOI: 10.1093/eurheartj/ehs079]
- Nalliah CJ**, Sanders P, Kottkamp H, Kalman JM. The role of obesity in atrial fibrillation. *Eur Heart J* 2015; pii: ehv486 [PMID: 26371114]
- Elahi MM**, Flatman S, Matata BM. Tracing the origins of postoperative atrial fibrillation: the concept of oxidative stress-mediated myocardial injury phenomenon. *Eur J Cardiovasc Prev Rehabil* 2008; **15**: 735-741 [PMID: 19020458 DOI: 10.1097/HJR.0b013e32831f738a]
- Zhang Q**, Liu T, Ng CY, Li G. Diabetes mellitus and atrial remodeling: mechanisms and potential upstream therapies. *Cardiovasc Ther* 2014; **32**: 233-241 [PMID: 25065462 DOI: 10.1111/1755-5922.12089]
- Abed HS**, Wittert GA. Obesity and atrial fibrillation. *Obes Rev* 2013; **14**: 929-938 [PMID: 23879190 DOI: 10.1111/obr.12056]
- Vargas-Uricoechea H**, Sierra-Torres CH. Thyroid hormones and the heart. *Horm Mol Biol Clin Investig* 2014; **18**: 15-26 [PMID: 25389997 DOI: 10.1515/hmbci-2013-0059]
- Goudis CA**, Korantzopoulos P, Ntalas IV, Kallergis EM, Ketikoglou DG. Obesity and atrial fibrillation: A comprehensive review of the pathophysiological mechanisms and links. *J Cardiol* 2015; **66**: 361-369 [PMID: 25959929 DOI: 10.1016/j.jjcc.2015.04.002]
- Goudis CA**, Korantzopoulos P, Ntalas IV, Kallergis EM, Liu T, Ketikoglou DG. Diabetes mellitus and atrial fibrillation: Pathophysiological mechanisms and potential upstream therapies. *Int J Cardiol* 2015; **184**: 617-622 [PMID: 25770841 DOI: 10.1016/j.ijcard.2015.03.052]
- De Caterina R**, Camm AJ. What is 'valvular' atrial fibrillation? A reappraisal. *Eur Heart J* 2014; **35**: 3328-3335 [PMID: 25265975 DOI: 10.1093/eurheartj/ehu352]
- Kumar KR**, Mandleywala SN, Link MS. Atrial and ventricular arrhythmias in hypertrophic cardiomyopathy. *Card Electrophysiol Clin* 2015; **7**: 173-186 [PMID: 26002384 DOI: 10.1016/j.ccep.2015.03.00]
- Riber LP**, Larsen TB, Christensen TD. Postoperative atrial fibrillation prophylaxis after lung surgery: systematic review and meta-analysis. *Ann Thorac Surg* 2014; **98**: 1989-1997 [PMID: 25283696 DOI: 10.1016/j.athoracsur.2014.06.069]
- Qaddoura A**, Kabali C, Drew D, van Oosten EM, Michael KA, Redfearn DP, Simpson CS, Baranchuk A. Obstructive sleep apnea as a predictor of atrial fibrillation after coronary artery bypass grafting: a systematic review and meta-analysis. *Can J Cardiol* 2014; **30**: 1516-1522 [PMID: 25475456 DOI: 10.1016/j.cjca.2014.10.014]
- Anumonwo JM**, Kalifa J. Risk factors and genetics of atrial fibrillation. *Cardiol Clin* 2014; **32**: 485-494 [PMID: 25443231 DOI: 10.1016/j.ccl.2014.07.007]
- Yadava M**, Hughey AB, Crawford TC. Postoperative atrial fibrillation: incidence, mechanisms, and clinical correlates. *Cardiol Clin* 2014; **32**: 627-636 [PMID: 25443241 DOI: 10.1016/j.ccl.2014.07.002]
- Akoum N**, Marrouche N. Assessment and impact of cardiac fibrosis on atrial fibrillation. *Curr Cardiol Rep* 2014; **16**: 518 [PMID: 24950676 DOI: 10.1007/s11886-014-0518]
- Berenfeld O**, Jalife J. Mechanisms of atrial fibrillation: rotors, ionic determinants, and excitation frequency. *Cardiol Clin* 2014; **32**: 495-506 [PMID: 25443232 DOI: 10.1016/j.ccl.2014.07.001]
- Heijman J**, Voigt N, Wehrens XH, Dobrev D. Calcium dysregulation in atrial fibrillation: the role of CaMKII. *Front Pharmacol* 2014; **5**: 30 [PMID: 24624086 DOI: 10.3389/fphar.2014.00030]
- Wolke C**, Bukowska A, Goette A, Lendeckel U. Redox control of cardiac remodeling in atrial fibrillation. *Biochim Biophys Acta* 2015; **1850**: 1555-1565 [PMID: 25513966 DOI: 10.1016/j.bbagen.2014.12.012]
- Jalife J**, Kaur K. Atrial remodeling, fibrosis, and atrial fibrillation. *Trends Cardiovasc Med* 2015; **25**: 475-484 [PMID: 25661032 DOI: 10.1016/j.tcm.2014.12.015]
- Corradi D**. Atrial fibrillation from the pathologist's perspective. *Cardiovasc Pathol* 2014; **23**: 71-84 [PMID: 24462196 DOI: 10.1016/j.carpath.2013.12.001]
- Dzeshka MS**, Lip GY, Snezhitskiy V, Shantsila E. Cardiac Fibrosis in Patients With Atrial Fibrillation: Mechanisms and Clinical Implications. *J Am Coll Cardiol* 2015; **66**: 943-959 [PMID: 26293766 DOI: 10.1016/j.jacc.2015.06.1313]
- Senoo K**, Lane D, Lip GY. Stroke and bleeding risk in atrial fibrillation. *Korean Circ J* 2014; **44**: 281-290 [PMID: 25278980 DOI: 10.4070/kcj.2014.44.5.281]
- Zimetbaum P**, Waks JW, Ellis ER, Glotzer TV, Passman RS. Role of atrial fibrillation burden in assessing thromboembolic risk. *Circ Arrhythm Electrophysiol* 2014; **7**: 1223-1229 [PMID: 25516580 DOI: 10.1161/CIRCEP.114.001356]
- Hirsh BJ**, Copeland-Halperin RS, Halperin JL. Fibrotic atrial cardiomyopathy, atrial fibrillation, and thromboembolism: mechanistic links and clinical inferences. *J Am Coll Cardiol* 2015; **65**: 2239-2251 [PMID: 25998669 DOI: 10.1016/j.jacc.2015.03.557]
- Hui DS**, Morley JE, Mikolajczak PC, Lee R. Atrial fibrillation: A major risk factor for cognitive decline. *Am Heart J* 2015; **169**: 448-456 [PMID: 25819850 DOI: 10.1016/j.ahj.2014.12.015]
- Jacobs V**, Cutler MJ, Day JD, Bunch TJ. Atrial fibrillation and dementia. *Trends Cardiovasc Med* 2015; **25**: 44-51 [PMID: 25442732]
- Luong C**, Barnes ME, Tsang TS. Atrial fibrillation and heart failure: cause or effect? *Curr Heart Fail Rep* 2014; **11**: 463-470 [PMID: 25258196 DOI: 10.1007/s11897-014-0229-1]
- Wijesurendra RS**, Casadei B. Atrial fibrillation: effects beyond the atrium? *Cardiovasc Res* 2015; **105**: 238-247 [PMID: 25587048 DOI: 10.1093/cvr/cvv001]
- Chen LY**, Benditt DG, Alonso A. Atrial fibrillation and its association with sudden cardiac death. *Circ J* 2014; **78**: 2588-2593 [PMID: 25262841]
- Potpara TS**, Lip GY. A brief history of 'lone' atrial fibrillation: from 'a peculiar pulse irregularity' to a modern public health concern. *Curr Pharm Des* 2015; **21**: 679-696 [PMID: 25269559]
- Andreassen L**, Nielsen JB, Olesen MS. Genetic aspects of lone atrial fibrillation: what do we know? *Curr Pharm Des* 2015; **21**: 667-678 [PMID: 25175087]
- Tello-Montoliu A**, Hernández-Romero D, Sanchez-Martínez M, Valdes M, Marín F. Lone atrial fibrillation - a diagnosis of exclusion. *Curr Pharm Des* 2015; **21**: 544-550 [PMID: 25175090]
- Andreassen L**, Nielsen JB, Christophersen IE, Holst AG, Sajadieh A, Tveit A, Haunsø S, Svendsen JH, Schmitt N, Olesen MS. Genetic modifier of the QTc interval associated with early-onset atrial fibrillation. *Can J Cardiol* 2013; **29**: 1234-1240 [PMID: 24074973 DOI: 10.1016/j.cjca.2013.06.009]
- Brugada R**, Tapscott T, Czerskiewicz GZ, Marian AJ, Iglesias A, Mont L, Brugada J, Girona J, Domingo A, Bachinski LL, Roberts R. Identification of a genetic locus for familial atrial fibrillation. *N Engl J Med* 1997; **336**: 905-911 [PMID: 9070470]

- 36 **Chen YH**, Xu SJ, Bendahhou S, Wang XL, Wang Y, Xu WY, Jin HW, Sun H, Su XY, Zhuang QN, Yang YQ, Li YB, Liu Y, Xu HJ, Li XF, Ma N, Mou CP, Chen Z, Barhanin J, Huang W. KCNQ1 gain-of-function mutation in familial atrial fibrillation. *Science* 2003; **299**: 251-254 [PMID: 12522251]
- 37 **Xia M**, Jin Q, Bendahhou S, He Y, Larroque MM, Chen Y, Zhou Q, Yang Y, Liu Y, Liu B, Zhu Q, Zhou Y, Lin J, Liang B, Li L, Dong X, Pan Z, Wang R, Wan H, Qiu W, Xu W, Eurlings P, Barhanin J, Chen Y. A Kir2.1 gain-of-function mutation underlies familial atrial fibrillation. *Biochem Biophys Res Commun* 2005; **332**: 1012-1019 [PMID: 15922306]
- 38 **Yang Y**, Xia M, Jin Q, Bendahhou S, Shi J, Chen Y, Liang B, Lin J, Liu Y, Liu B, Zhou Q, Zhang D, Wang R, Ma N, Su X, Niu K, Pei Y, Xu W, Chen Z, Wan H, Cui J, Barhanin J, Chen Y. Identification of a KCNE2 gain-of-function mutation in patients with familial atrial fibrillation. *Am J Hum Genet* 2004; **75**: 899-905 [PMID: 15368194]
- 39 **Berenfeld O**, Zaitsev AV, Mironov SF, Pertsov AM, Jalife J. Frequency-dependent breakdown of wave propagation into fibrillatory conduction across the pectinate muscle network in the isolated sheep right atrium. *Circ Res* 2002; **90**: 1173-1180 [PMID: 12065320]
- 40 **Chen J**, Mandapati R, Berenfeld O, Skanes AC, Gray RA, Jalife J. Dynamics of wavelets and their role in atrial fibrillation in the isolated sheep heart. *Cardiovasc Res* 2000; **48**: 220-232 [PMID: 11054469]
- 41 **Mandapati R**, Skanes A, Chen J, Berenfeld O, Jalife J. Stable microreentrant sources as a mechanism of atrial fibrillation in the isolated sheep heart. *Circulation* 2000; **101**: 194-199 [PMID: 10637208]
- 42 **Skanes AC**, Mandapati R, Berenfeld O, Davidenko JM, Jalife J. Spatiotemporal periodicity during atrial fibrillation in the isolated sheep heart. *Circulation* 1998; **98**: 1236-1248 [PMID: 9743516]
- 43 **Tanaka K**, Zlochiver S, Vikstrom KL, Yamazaki M, Moreno J, Klos M, Zaitsev AV, Vaidyanathan R, Auerbach DS, Landas S, Guiraudon G, Jalife J, Berenfeld O, Kalifa J. Spatial distribution of fibrosis governs fibrillation wave dynamics in the posterior left atrium during heart failure. *Circ Res* 2007; **101**: 839-847 [PMID: 17704207]
- 44 **Gudbjartsson DF**, Arnar DO, Helgadóttir A, Gretarsdóttir S, Holm H, Sigurdsson A, Jonasdóttir A, Baker A, Thorleifsson G, Kristjansson K, Palsson A, Blondal T, Sulem P, Backman VM, Hardarson GA, Palsdóttir E, Helgason A, Sigurjonsdóttir R, Sverrisson JT, Kostulas K, Ng MC, Baum L, So WY, Wong KS, Chan JC, Furie KL, Greenberg SM, Sale M, Kelly P, MacRae CA, Smith EE, Rosand J, Hillert J, Ma RC, Ellinor PT, Thorgeirsson G, Gulcher JR, Kong A, Thorsteinsdóttir U, Stefansson K. Variants conferring risk of atrial fibrillation on chromosome 4q25. *Nature* 2007; **448**: 353-357 [PMID: 17603472]
- 45 **Benjamin EJ**, Rice KM, Arking DE, Pfeuffer A, van Noord C, Smith AV, Schnabel RB, Bis JC, Boerwinkle E, Sinner MF, Dehghan A, Lubitz SA, D'Agostino RB, Lumley T, Ehret GB, Heeringa J, Aspelund T, Newton-Cheh C, Larson MG, Marcante KD, Soliman EZ, Rivadeneira F, Wang TJ, Eiriksdóttir G, Levy D, Psaty BM, Li M, Chamberlain AM, Hofman A, Vasan RS, Harris TB, Rotter JI, Kao WH, Agarwal SK, Stricker BH, Wang K, Launer LJ, Smith NL, Chakravarti A, Uitterlinden AG, Wolf PA, Sotoodehnia N, Köttgen A, van Duijn CM, Meitinger T, Mueller M, Perz S, Steinbeck G, Wichmann HE, Lunetta KL, Heckbert SR, Gudnason V, Alonso A, Kääb S, Ellinor PT, Witteman JC. Variants in ZFX3 are associated with atrial fibrillation in individuals of European ancestry. *Nat Genet* 2009; **41**: 879-881 [PMID: 19597492 DOI: 10.1038/ng.416]
- 46 **Ellinor PT**, Lunetta KL, Albert CM, Glazer NL, Ritchie MD, Smith AV, Arking DE, Müller-Nurasyid M, Krijthe BP, Lubitz SA, Bis JC, Chung MK, Dörr M, Ozaki K, Roberts JD, Smith JG, Pfeuffer A, Sinner MF, Lohman K, Ding J, Smith NL, Smith JD, Rienstra M, Rice KM, Van Wagoner DR, Magnani JW, Wakili R, Clauss S, Rotter JI, Steinbeck G, Launer LJ, Davies RW, Borkovich M, Harris TB, Lin H, Völker U, Völzke H, Milan DJ, Hofman A, Boerwinkle E, Chen LY, Soliman EZ, Voight BF, Li G, Chakravarti A, Kubo M, Tedrow UB, Rose LM, Ridker PM, Conen D, Tsunoda T, Furukawa T, Sotoodehnia N, Xu S, Kamatani N, Levy D, Nakamura Y, Parvez B, Mahida S, Furie KL, Rosand J, Muhammad R, Psaty BM, Meitinger T, Perz S, Wichmann HE, Witteman JC, Kao WH, Kathiresan S, Roden DM, Uitterlinden AG, Rivadeneira F, McKnight B, Sjögren M, Newman AB, Liu Y, Gollob MH, Melander O, Tanaka T, Stricker BH, Felix SB, Alonso A, Darbar D, Barnard J, Chasman DI, Heckbert SR, Benjamin EJ, Gudnason V, Kääb S. Meta-analysis identifies six new susceptibility loci for atrial fibrillation. *Nat Genet* 2012; **44**: 670-675 [PMID: 22544366 DOI: 10.1038/ng.2261]
- 47 **Ellinor PT**, Lunetta KL, Glazer NL, Pfeuffer A, Alonso A, Chung MK, Sinner MF, de Bakker PI, Mueller M, Lubitz SA, Fox E, Darbar D, Smith NL, Smith JD, Schnabel RB, Soliman EZ, Rice KM, Van Wagoner DR, Beckmann BM, van Noord C, Wang K, Ehret GB, Rotter JI, Hazen SL, Steinbeck G, Smith AV, Launer LJ, Harris TB, Makino S, Nelis M, Milan DJ, Perz S, Esko T, Köttgen A, Moebus S, Newton-Cheh C, Li M, Möhlenkamp S, Wang TJ, Kao WH, Vasan RS, Nöthen MM, MacRae CA, Stricker BH, Hofman A, Uitterlinden AG, Levy D, Boerwinkle E, Metspalu A, Topol EJ, Chakravarti A, Gudnason V, Psaty BM, Roden DM, Meitinger T, Wichmann HE, Witteman JC, Barnard J, Arking DE, Benjamin EJ, Heckbert SR, Kääb S. Common variants in KCNN3 are associated with lone atrial fibrillation. *Nat Genet* 2010; **42**: 240-244 [PMID: 20173747 DOI: 10.1038/ng.537]
- 48 **Gudbjartsson DF**, Holm H, Gretarsdóttir S, Thorleifsson G, Walters GB, Thorgeirsson G, Gulcher J, Mathiesen EB, Njølstad I, Nyrnes A, Wilsgaard T, Hald EM, Hveem K, Stoltenberg C, Kucera G, Stubblefield T, Carter S, Roden D, Ng MC, Baum L, So WY, Wong KS, Chan JC, Gieger C, Wichmann HE, Gschwendtner A, Dichgans M, Kühlenbäumer G, Berger K, Ringelstein EB, Bevan S, Markus HS, Kostulas K, Hillert J, Sveinbjörnsdóttir S, Valdimarsson EM, Løchen ML, Ma RC, Darbar D, Kong A, Arnar DO, Thorsteinsdóttir U, Stefansson K. A sequence variant in ZFX3 on 16q22 associates with atrial fibrillation and ischemic stroke. *Nat Genet* 2009; **41**: 876-878 [PMID: 19597491 DOI: 10.1038/ng.417]
- 49 **Schnabel RB**, Kerr KF, Lubitz SA, Alkylbekova EL, Marcus GM, Sinner MF, Magnani JW, Wolf PA, Deo R, Lloyd-Jones DM, Lunetta KL, Mehra R, Levy D, Fox ER, Arking DE, Mosley TH, Müller-Nurasyid M, Young TR, Wichmann HE, Seshadri S, Farlow DN, Rotter JI, Soliman EZ, Glazer NL, Wilson JG, Breteler MM, Sotoodehnia N, Newton-Cheh C, Kääb S, Ellinor PT, Alonso A, Benjamin EJ, Heckbert SR. Large-scale candidate gene analysis in whites and African Americans identifies IL6R polymorphism in relation to atrial fibrillation: the National Heart, Lung, and Blood Institute's Candidate Gene Association Resource (CARE) project. *Circ Cardiovasc Genet* 2011; **4**: 557-564 [PMID: 21846873 DOI: 10.1161/CIRCGENETICS.110.959197]
- 50 **Amin AS**, Tan HL, Wilde AA. Cardiac ion channels in health and disease. *Heart Rhythm* 2010; **7**: 117-126 [PMID: 19875343 DOI: 10.1016/j.hrthm.2009.08.005]
- 51 **Amin AS**, Bhuiyan ZA. SCN5A mutations in atrial fibrillation. *Heart Rhythm* 2010; **7**: 1870-1871 [PMID: 20850563 DOI: 10.1016/j.hrthm.2010.09.012]
- 52 **Blana A**, Kaese S, Fortmüller L, Laakmann S, Damke D, van Bragt K, Eckstein J, Piccini I, Kirchhefer U, Nattel S, Breithardt G, Carmeliet P, Carmeliet E, Schotten U, Verheule S, Kirchhof P, Fabritz L. Knock-in gain-of-function sodium channel mutation prolongs atrial action potentials and alters atrial vulnerability. *Heart Rhythm* 2010; **7**: 1862-1869 [PMID: 20728579 DOI: 10.1016/j.hrthm.2010.08.016]
- 53 **Chen S**, Wang C, Wang X, Xu C, Wu M, Wang P, Tu X, Wang QK. Significant Association Between CAV1 Variant rs3807989 on 7p31 and Atrial Fibrillation in a Chinese Han Population. *J Am Heart Assoc* 2015; **4**: pii: e001980 [PMID: 25953654 DOI: 10.1161/JAHA.115.001980]
- 54 **Laitinen-Forsblom PJ**, Mäkinen P, Mäkinen H, Yli-Mäyry S, Virtanen V, Kontula K, Aalto-Setälä K. SCN5A mutation

- associated with cardiac conduction defect and atrial arrhythmias. *J Cardiovasc Electrophysiol* 2006; **17**: 480-485 [PMID: 16684018]
- 55 **Li Q**, Huang H, Liu G, Lam K, Rutberg J, Green MS, Birnie DH, Lemery R, Chahine M, Gollob MH. Gain-of-function mutation of Nav1.5 in atrial fibrillation enhances cellular excitability and lowers the threshold for action potential firing. *Biochem Biophys Res Commun* 2009; **380**: 132-137 [PMID: 19167345 DOI: 10.1016/j.bbrc.2009.01.052]
- 56 **Makiyama T**, Akao M, Shizuta S, Doi T, Nishiyama K, Oka Y, Ohno S, Nishio Y, Tsuji K, Itoh H, Kimura T, Kita T, Horie M. A novel SCN5A gain-of-function mutation M1875T associated with familial atrial fibrillation. *J Am Coll Cardiol* 2008; **52**: 1326-1334 [PMID: 1829244 DOI: 10.1016/j.jacc.2008.07.013]
- 57 **Benito B**, Brugada R, Perich RM, Lizotte E, Cinca J, Mont L, Berrueto A, Tolosana JM, Freixa X, Brugada P, Brugada J. A mutation in the sodium channel is responsible for the association of long QT syndrome and familial atrial fibrillation. *Heart Rhythm* 2008; **5**: 1434-1440 [PMID: 18929331 DOI: 10.1016/j.hrthm.2008.07.013]
- 58 **Calloe K**, Schmitt N, Grubb S, Pfeiffer R, David JP, Kanter R, Cordeiro JM, Antzelevitch C. Multiple arrhythmic syndromes in a newborn, owing to a novel mutation in SCN5A. *Can J Physiol Pharmacol* 2011; **89**: 723-736 [PMID: 21895525 DOI: 10.1139/y11-070]
- 59 **Dolz-Gaitón P**, Núñez M, Núñez L, Barana A, Amorós I, Matamoros M, Pérez-Hernández M, González de la Fuente M, Alvarez-López M, Macías-Ruiz R, Tercedor-Sánchez L, Jiménez-Jáimez J, Delpón E, Caballero R, Tamargo J. Functional characterization of a novel frameshift mutation in the C-terminus of the Nav1.5 channel underlying a Brugada syndrome with variable expression in a Spanish family. *PLoS One* 2013; **8**: e81493 [PMID: 24363796 DOI: 10.1371/journal.pone.0081493]
- 60 **Olson TM**, Michels VV, Ballew JD, Reyna SP, Karst ML, Herron KJ, Horton SC, Rodeheffer RJ, Anderson JL. Sodium channel mutations and susceptibility to heart failure and atrial fibrillation. *JAMA* 2005; **293**: 447-454 [PMID: 15671429]
- 61 **Ziyadeh-Isleem A**, Clatot J, Duchatelet S, Gandjbakhch E, Denjoy I, Hidden-Lucet F, Hatem S, Deschênes I, Coulombe A, Neyroud N, Guicheney P. A truncating SCN5A mutation combined with genetic variability causes sick sinus syndrome and early atrial fibrillation. *Heart Rhythm* 2014; **11**: 1015-1023 [PMID: 24582607 DOI: 10.1016/j.hrthm.2014.02.021]
- 62 **Rossenbacker T**, Carroll SJ, Liu H, Kuipéri C, de Ravel TJ, Devriendt K, Carmeliet P, Kass RS, Heidebüchel H. Novel pore mutation in SCN5A manifests as a spectrum of phenotypes ranging from atrial flutter, conduction disease, and Brugada syndrome to sudden cardiac death. *Heart Rhythm* 2004; **1**: 610-615 [PMID: 15851228]
- 63 **Watanabe H**, Darbar D, Kaiser DW, Jiramongkolchai K, Chopra S, Donahue BS, Kannankeril PJ, Roden DM. Mutations in sodium channel β 1- and β 2-subunits associated with atrial fibrillation. *Circ Arrhythm Electrophysiol* 2009; **2**: 268-275 [PMID: 19808477 DOI: 10.1161/CIRCEP.108.779181]
- 64 **Wang P**, Yang Q, Wu X, Yang Y, Shi L, Wang C, Wu G, Xia Y, Yang B, Zhang R, Xu C, Cheng X, Li S, Zhao Y, Fu F, Liao Y, Fang F, Chen Q, Tu X, Wang QK. Functional dominant-negative mutation of sodium channel subunit gene SCN3B associated with atrial fibrillation in a Chinese GenID population. *Biochem Biophys Res Commun* 2010; **398**: 98-104 [PMID: 20558140 DOI: 10.1016/j.bbrc.2010.06.042]
- 65 **Olesen MS**, Jespersen T, Nielsen JB, Liang B, Møller DV, Hedley P, Christiansen M, Varró A, Olesen SP, Haunsø S, Schmitt N, Svendsen JH. Mutations in sodium channel β -subunit SCN3B are associated with early-onset lone atrial fibrillation. *Cardiovasc Res* 2011; **89**: 786-793 [PMID: 21051419 DOI: 10.1093/cvr/cvq348]
- 66 **Li RG**, Wang Q, Xu YJ, Zhang M, Qu XK, Liu X, Fang WY, Yang YQ. Mutations of the SCN4B-encoded sodium channel β 4 subunit in familial atrial fibrillation. *Int J Mol Med* 2013; **32**: 144-150 [PMID: 23604097 DOI: 10.3892/ijmm.2013.1355]
- 67 **Olesen MS**, Holst AG, Svendsen JH, Haunsø S, Tfelt-Hansen J, SCN1Bb R214Q found in 3 patients: 1 with Brugada syndrome and 2 with lone atrial fibrillation. *Heart Rhythm* 2012; **9**: 770-773 [PMID: 22155598 DOI: 10.1016/j.hrthm.2011.12.005]
- 68 **Remme CA**, Wilde AA, Bezzina CR. Cardiac sodium channel overlap syndromes: different faces of SCN5A mutations. *Trends Cardiovasc Med* 2008; **18**: 78-87 [PMID: 18436145 DOI: 10.1016/j.tcm.2008.01.002]
- 69 **Rook MB**, Evers MM, Vos MA, Bierhuizen MF. Biology of cardiac sodium channel Nav1.5 expression. *Cardiovasc Res* 2012; **93**: 12-23 [PMID: 21937582 DOI: 10.1093/cvr/cvr252]
- 70 **Veerman CC**, Wilde AA, Lodder EM. The cardiac sodium channel gene SCN5A and its gene product Nav1.5: Role in physiology and pathophysiology. *Gene* 2015; **573**: 177-187 [PMID: 26361848 DOI: 10.1016/j.gene.2015.08.062]
- 71 **Macri V**, Mahida SN, Zhang ML, Sinner MF, Dolmatova EV, Tucker NR, McLellan M, Shea MA, Milan DJ, Lunetta KL, Benjamin EJ, Ellinor PT. A novel trafficking-defective HCN4 mutation is associated with early-onset atrial fibrillation. *Heart Rhythm* 2014; **11**: 1055-1062 [PMID: 24607718 DOI: 10.1016/j.hrthm.2014.03.002]
- 72 **Hong K**, Piper DR, Diaz-Valdecantos A, Brugada J, Oliva A, Burashnikov E, Santos-de-Soto J, Gueso-Montero J, Diaz-Enfante E, Brugada P, Sachse F, Sanguinetti MC, Brugada R. De novo KCNQ1 mutation responsible for atrial fibrillation and short QT syndrome in utero. *Cardiovasc Res* 2005; **68**: 433-440 [PMID: 16109388]
- 73 **Lundby A**, Ravn LS, Svendsen JH, Olesen SP, Schmitt N. KCNQ1 mutation Q147R is associated with atrial fibrillation and prolonged QT interval. *Heart Rhythm* 2007; **4**: 1532-1541 [PMID: 17997361]
- 74 **Kharche S**, Adeniran I, Stott J, Law P, Boyett MR, Hancox JC, Zhang H. Pro-arrhythmogenic effects of the S140G KCNQ1 mutation in human atrial fibrillation - insights from modelling. *J Physiol* 2012; **590**: 4501-4514 [PMID: 22508963 DOI: 10.1113/jphysiol.2012.229146]
- 75 **Das S**, Makino S, Melman YF, Shea MA, Goyal SB, Rosenzweig A, Macrae CA, Ellinor PT. Mutation in the S3 segment of KCNQ1 results in familial lone atrial fibrillation. *Heart Rhythm* 2009; **6**: 1146-1153 [PMID: 19632626 DOI: 10.1016/j.hrthm.2009.04.015]
- 76 **El Harchi A**, Zhang H, Hancox JC. The S140G KCNQ1 atrial fibrillation mutation affects 'I(KS)' profile during both atrial and ventricular action potentials. *J Physiol Pharmacol* 2010; **61**: 759-764 [PMID: 21224508]
- 77 **Bartos DC**, Duchatelet S, Burgess DE, Klug D, Denjoy I, Peat R, Lupoglazoff JM, Fressart V, Berthet M, Ackerman MJ, January CT, Guicheney P, Delisle BP. R231C mutation in KCNQ1 causes long QT syndrome type 1 and familial atrial fibrillation. *Heart Rhythm* 2011; **8**: 48-55 [PMID: 20850564 DOI: 10.1016/j.hrthm.2010.09.010]
- 78 **Hasegawa K**, Ohno S, Ashihara T, Itoh H, Ding WG, Toyoda F, Makiyama T, Aoki H, Nakamura Y, Delisle BP, Matsuura H, Horie M. A novel KCNQ1 missense mutation identified in a patient with juvenile-onset atrial fibrillation causes constitutively open IKs channels. *Heart Rhythm* 2014; **11**: 67-75 [PMID: 24096004 DOI: 10.1016/j.hrthm.2013.09.073]
- 79 **Ki CS**, Jung CL, Kim HJ, Baek KH, Park SJ, On YK, Kim KS, Noh SJ, Youm JB, Kim JS, Cho H. A KCNQ1 mutation causes age-dependant bradycardia and persistent atrial fibrillation. *Pflugers Arch* 2014; **466**: 529-540 [PMID: 23989646 DOI: 10.1007/s00424-013-1337-6]
- 80 **Ravn LS**, Aizawa Y, Pollevick GD, Hofman-Bang J, Cordeiro JM, Diken U, Jensen G, Wu Y, Burashnikov E, Haunsø S, Guerchicoff A, Hu D, Svendsen JH, Christiansen M, Antzelevitch C. Gain of function in IKs secondary to a mutation in KCNE5 associated with atrial fibrillation. *Heart Rhythm* 2008; **5**: 427-435 [PMID: 18313602 DOI: 10.1016/j.hrthm.2007.12.019]
- 81 **Nielsen JB**, Bentzen BH, Olesen MS, David JP, Olesen SP, Haunsø S, Svendsen JH, Schmitt N. Gain-of-function mutations in potassium channel subunit KCNE2 associated with early-onset lone atrial fibrillation. *Biomark Med* 2014; **8**: 557-570 [PMID: 24796621 DOI: 10.2217/bmm.13.137]

- 82 **Deo M**, Ruan Y, Pandit SV, Shah K, Berenfeld O, Blafox A, Cerrone M, Noujaim SF, Denegri M, Jalife J, Priori SG. KCNJ2 mutation in short QT syndrome 3 results in atrial fibrillation and ventricular proarrhythmia. *Proc Natl Acad Sci USA* 2013; **110**: 4291-4296 [PMID: 23440193 DOI: 10.1073/pnas.1218154110]
- 83 **Kharache S**, Garratt CJ, Boyett MR, Inada S, Holden AV, Hancox JC, Zhang H. Atrial proarrhythmia due to increased inward rectifier current (I(K1)) arising from KCNJ2 mutation--a simulation study. *Prog Biophys Mol Biol* 2008; **98**: 186-197 [PMID: 19041665 DOI: 10.1016/j.pbiomolbio.2008.10.010]
- 84 **Villafañe J**, Fischbach P, Gebauer R. Short QT syndrome manifesting with neonatal atrial fibrillation and bradycardia. *Cardiology* 2014; **128**: 236-240 [PMID: 24818999 DOI: 10.1159/000360758]
- 85 **Chelu MG**, Sarma S, Sood S, Wang S, van Oort RJ, Skapura DG, Li N, Santonastasi M, Müller FU, Schmitz W, Schotten U, Anderson ME, Valderrábano M, Dobrev D, Wehrens XH. Calmodulin kinase II-mediated sarcoplasmic reticulum Ca²⁺ leak promotes atrial fibrillation in mice. *J Clin Invest* 2009; **119**: 1940-1951 [PMID: 19603549]
- 86 **Shan J**, Xie W, Betzenhauser M, Reiken S, Chen BX, Wronska A, Marks AR. Calcium leak through ryanodine receptors leads to atrial fibrillation in 3 mouse models of catecholaminergic polymorphic ventricular tachycardia. *Circ Res* 2012; **111**: 708-717 [PMID: 22828895 DOI: 10.1161/CIRCRESAHA.112.273342]
- 87 **Li N**, Wang T, Wang W, Cutler MJ, Wang Q, Voigt N, Rosenbaum DS, Dobrev D, Wehrens XH. Inhibition of CaMKII phosphorylation of RyR2 prevents induction of atrial fibrillation in FKBP12.6 knockout mice. *Circ Res* 2012; **110**: 465-470 [PMID: 22158709 DOI: 10.1161/CIRCRESAHA.111.253229]
- 88 **Zhang Y**, Matthews GD, Lei M, Huang CL. Abnormal Ca(2+) homeostasis, atrial arrhythmogenesis, and sinus node dysfunction in murine hearts modeling RyR2 modification. *Front Physiol* 2013; **4**: 150 [PMID: 23805105 DOI: 10.3389/fphys.2013.00150]
- 89 **Beavers DL**, Wang W, Ather S, Voigt N, Garbino A, Dixit SS, Landstrom AP, Li N, Wang Q, Olivetto I, Dobrev D, Ackerman MJ, Wehrens XH. Mutation E169K in junctophilin-2 causes atrial fibrillation due to impaired RyR2 stabilization. *J Am Coll Cardiol* 2013; **62**: 2010-2019 [PMID: 23973696 DOI: 10.1016/j.jacc.2013.06.052]
- 90 **Chiang DY**, Li N, Wang Q, Alsina KM, Quick AP, Reynolds JO, Wang G, Skapura D, Voigt N, Dobrev D, Wehrens XH. Impaired local regulation of ryanodine receptor type 2 by protein phosphatase 1 promotes atrial fibrillation. *Cardiovasc Res* 2014; **103**: 178-187 [PMID: 24812280 DOI: 10.1093/cvr/cvu123]
- 91 **Di Pino A**, Caruso E, Costanzo L, Guccione P. A novel RyR2 mutation in a 2-year-old baby presenting with atrial fibrillation, atrial flutter, and atrial ectopic tachycardia. *Heart Rhythm* 2014; **11**: 1480-1483 [PMID: 24793461 DOI: 10.1016/j.hrthm.2014.04.037]
- 92 **Zhabiyev P**, Hiess F, Wang R, Liu Y, Wayne Chen SR, Oudit GY. S4153R is a gain-of-function mutation in the cardiac Ca(2+) release channel ryanodine receptor associated with catecholaminergic polymorphic ventricular tachycardia and paroxysmal atrial fibrillation. *Can J Cardiol* 2013; **29**: 993-996 [PMID: 23498838 DOI: 10.1016/j.cjca.2012.12.019]
- 93 **Kazemian P**, Gollob MH, Pantano A, Oudit GY. A novel mutation in the RYR2 gene leading to catecholaminergic polymorphic ventricular tachycardia and paroxysmal atrial fibrillation: dose-dependent arrhythmia-event suppression by β -blocker therapy. *Can J Cardiol* 2011; **27**: 870.e7-870.e10 [PMID: 21652165 DOI: 10.1016/j.cjca.2011.02.003]
- 94 **King JH**, Wickramarachchi C, Kua K, Du Y, Jeevaratnam K, Matthews HR, Grace AA, Huang CL, Fraser JA. Loss of Nav1.5 expression and function in murine atria containing the RyR2-P2328S gain-of-function mutation. *Cardiovasc Res* 2013; **99**: 751-759 [PMID: 23723061 DOI: 10.1093/cvr/cvt141]
- 95 **Zhang Y**, Fraser JA, Jeevaratnam K, Hao X, Hothi SS, Grace AA, Lei M, Huang CL. Acute atrial arrhythmogenicity and altered Ca(2+) homeostasis in murine RyR2-P2328S hearts. *Cardiovasc Res* 2011; **89**: 794-804 [PMID: 20621925 DOI: 10.1093/cvr/cvq229]
- 96 **Thibodeau IL**, Xu J, Li Q, Liu G, Lam K, Veinot JP, Birnie DH, Jones DL, Krahn AD, Lemery R, Nicholson BJ, Gollob MH. Paradigm of genetic mosaicism and lone atrial fibrillation: physiological characterization of a connexin 43-deletion mutant identified from atrial tissue. *Circulation* 2010; **122**: 236-244 [PMID: 20606116 DOI: 10.1161/CIRCULATIONAHA.110.961227]
- 97 **Tuomi JM**, Tyml K, Jones DL. Atrial tachycardia/fibrillation in the connexin 43 G60S mutant (Oculodentodigital dysplasia) mouse. *Am J Physiol Heart Circ Physiol* 2011; **300**: H1402-H1411 [PMID: 21239638 DOI: 10.1152/ajpheart.01094.2010]
- 98 **Delmar M**, Makita N. Cardiac connexins, mutations and arrhythmias. *Curr Opin Cardiol* 2012; **27**: 236-241 [PMID: 22382502 DOI: 10.1097/HCO.0b013e328352220e]
- 99 **Gollob MH**, Jones DL, Krahn AD, Danis L, Gong XQ, Shao Q, Liu X, Veinot JP, Tang AS, Stewart AF, Tesson F, Klein GJ, Yee R, Skanes AC, Guiraudon GM, Ebihara L, Bai D. Somatic mutations in the connexin 40 gene (GJA5) in atrial fibrillation. *N Engl J Med* 2006; **354**: 2677-2688 [PMID: 16790700]
- 100 **Yang YQ**, Liu X, Zhang XL, Wang XH, Tan HW, Shi HF, Jiang WF, Fang WY. Novel connexin40 missense mutations in patients with familial atrial fibrillation. *Europace* 2010; **12**: 1421-1427 [PMID: 20650941 DOI: 10.1093/europace/euq274]
- 101 **Yang YQ**, Zhang XL, Wang XH, Tan HW, Shi HF, Jiang WF, Fang WY, Liu X. Connexin40 nonsense mutation in familial atrial fibrillation. *Int J Mol Med* 2010; **26**: 605-610 [PMID: 20818502]
- 102 **Christophersen IE**, Holmegard HN, Jabbari J, Sajadieh A, Haunsø S, Tveit A, Svendsen JH, Olesen MS. Rare variants in GJA5 are associated with early-onset lone atrial fibrillation. *Can J Cardiol* 2013; **29**: 111-116 [PMID: 23040431 DOI: 10.1016/j.cjca.2012.08.002]
- 103 **Shi HF**, Yang JF, Wang Q, Li RG, Xu YJ, Qu XK, Fang WY, Liu X, Yang YQ. Prevalence and spectrum of GJA5 mutations associated with lone atrial fibrillation. *Mol Med Rep* 2013; **7**: 767-774 [PMID: 23292621 DOI: 10.3892/mmr.2012.1252]
- 104 **Sun Y**, Yang YQ, Gong XQ, Wang XH, Li RG, Tan HW, Liu X, Fang WY, Bai D. Novel germline GJA5/connexin40 mutations associated with lone atrial fibrillation impair gap junctional intercellular communication. *Hum Mutat* 2013; **34**: 603-609 [PMID: 23348765 DOI: 10.1002/humu.22278]
- 105 **Roberts JD**, Longoria J, Poon A, Gollob MH, Dewland TA, Kwok PY, Olgin JE, Deo RC, Marcus GM. Targeted deep sequencing reveals no definitive evidence for somatic mosaicism in atrial fibrillation. *Circ Cardiovasc Genet* 2015; **8**: 50-57 [PMID: 25406240 DOI: 10.1161/CIRCGENETICS.114.000650]
- 106 **Gemel J**, Simon AR, Patel D, Xu Q, Matiukas A, Veenstra RD, Beyer EC. Degradation of a connexin40 mutant linked to atrial fibrillation is accelerated. *J Mol Cell Cardiol* 2014; **74**: 330-339 [PMID: 24973497 DOI: 10.1016/j.yjmcc.2014.06.010]
- 107 **Sun Y**, Hills MD, Ye WG, Tong X, Bai D. Atrial fibrillation-linked germline GJA5/connexin40 mutants showed an increased hemichannel function. *PLoS One* 2014; **9**: e95125 [PMID: 24733048 DOI: 10.1371/journal.pone.0095125]
- 108 **Sun Y**, Tong X, Chen H, Huang T, Shao Q, Huang W, Laird DW, Bai D. An atrial-fibrillation-linked connexin40 mutant is retained in the endoplasmic reticulum and impairs the function of atrial gap-junction channels. *Dis Model Mech* 2014; **7**: 561-569 [PMID: 24626989 DOI: 10.1242/dmm.013813]
- 109 **Patel D**, Gemel J, Xu Q, Simon AR, Lin X, Matiukas A, Beyer EC, Veenstra RD. Atrial fibrillation-associated connexin40 mutants make hemichannels and synergistically form gap junction channels with novel properties. *FEBS Lett* 2014; **588**: 1458-1464 [PMID: 24457199 DOI: 10.1016/j.febslet.2014.01.010]
- 110 **Bai D**. Atrial fibrillation-linked GJA5/connexin40 mutants impaired gap junctions via different mechanisms. *FEBS Lett* 2014; **588**: 1238-1243 [PMID: 24656738 DOI: 10.1016/j.febslet.2014.02.064]
- 111 **Lübke-meier I**, Andrié R, Lickfett L, Bosen F, Stöckigt F, Dobrowolski R, Draffehn AM, Fregeac J, Bukauskas FF, Schrickel JW, Willecke K. The Connexin40A96S mutation from a patient with atrial fibrillation causes decreased atrial conduction velocities and sustained episodes of induced atrial

- fibrillation in mice. *J Mol Cell Cardiol* 2013; **65**: 19-32 [PMID: 24060583 DOI: 10.1016/j.jmcc.2013.09.008]
- 112 **Postma AV**, van de Meerakker JB, Mathijssen IB, Barnett P, Christoffels VM, Ilgun A, Lam J, Wilde AA, Lekanne Deprez RH, Moorman AF. A gain-of-function TBX5 mutation is associated with atypical Holt-Oram syndrome and paroxysmal atrial fibrillation. *Circ Res* 2008; **102**: 1433-1442 [PMID: 18451335 DOI: 10.1161/CIRCRESAHA.107.168294]
- 113 **Gutierrez-Roelens I**, De Roy L, Ovaert C, Sluysmans T, Devriendt K, Brunner HG, Viskula M. A novel CSX/NKX2-5 mutation causes autosomal-dominant AV block: are atrial fibrillation and syncope part of the phenotype? *Eur J Hum Genet* 2006; **14**: 1313-1316 [PMID: 16896344]
- 114 **Huang RT**, Xue S, Xu YJ, Zhou M, Yang YQ. A novel NKX2.5 loss-of-function mutation responsible for familial atrial fibrillation. *Int J Mol Med* 2013; **31**: 1119-1126 [PMID: 23525379 DOI: 10.3892/ijmm.2013.1316]
- 115 **Xie WH**, Chang C, Xu YJ, Li RG, Qu XK, Fang WY, Liu X, Yang YQ. Prevalence and spectrum of Nkx2.5 mutations associated with idiopathic atrial fibrillation. *Clinics (Sao Paulo)* 2013; **68**: 777-784 [PMID: 23778487 DOI: 10.6061/clinics/2013(06)09]
- 116 **Yu H**, Xu JH, Song HM, Zhao L, Xu WJ, Wang J, Li RG, Xu L, Jiang WF, Qiu XB, Jiang JQ, Qu XK, Liu X, Fang WY, Jiang JF, Yang YQ. Mutational spectrum of the NKX2-5 gene in patients with lone atrial fibrillation. *Int J Med Sci* 2014; **11**: 554-563 [PMID: 24782644 DOI: 10.7150/ijms.8407]
- 117 **Yuan F**, Qiu XB, Li RG, Qu XK, Wang J, Xu YJ, Liu X, Fang WY, Yang YQ, Liao DN. A novel NKX2-5 loss-of-function mutation predisposes to familial dilated cardiomyopathy and arrhythmias. *Int J Mol Med* 2015; **35**: 478-486 [PMID: 25503402 DOI: 10.3892/ijmm.2014.2029]
- 118 **Huang J**, Zhang DF, Sun YM, Li RG, Qiu XB, Qu XK, Liu X, Fang WY, Yang YQ. NKX2-6 mutation predisposes to familial atrial fibrillation. *Int J Mol Med* 2014; **34**: 1581-1590 [PMID: 25319568 DOI: 10.3892/ijmm.2014.1971]
- 119 **Zhang X**, Chen S, Yoo S, Chakrabarti S, Zhang T, Ke T, Oberti C, Yong SL, Fang F, Li L, de la Fuente R, Wang L, Chen Q, Wang QK. Mutation in nuclear pore component NUP155 leads to atrial fibrillation and early sudden cardiac death. *Cell* 2008; **135**: 1017-1027 [PMID: 19070573 DOI: 10.1016/j.cell.2008.10.022]
- 120 **Hua R**, MacLeod SL, Polina I, Moghtadaei M, Jansen HJ, Bogachev O, O'Blens SB, Sapp JL, Legare JF, Rose RA. Effects of Wild-Type and Mutant Forms of Atrial Natriuretic Peptide on Atrial Electrophysiology and Arrhythmogenesis. *Circ Arrhythm Electrophysiol* 2015; **8**: 1240-1254 [PMID: 26227000]
- 121 **Li QY**, Newbury-Ecob RA, Terrett JA, Wilson DI, Curtis AR, Yi CH, Gebuhr T, Bullen PJ, Robson SC, Strachan T, Bonnet D, Lyonnet S, Young ID, Raeburn JA, Buckler AJ, Law DJ, Brook JD. Holt-Oram syndrome is caused by mutations in TBX5, a member of the Brachyury (T) gene family. *Nat Genet* 1997; **15**: 21-29 [PMID: 8988164]
- 122 **Costa MW**, Guo G, Wolstein O, Vale M, Castro ML, Wang L, Otway R, Riek P, Cochrane N, Furtado M, Semsarian C, Weintraub RG, Yeoh T, Hayward C, Keogh A, Macdonald P, Feneley M, Graham RM, Seidman JG, Seidman CE, Rosenthal N, Fatkin D, Harvey RP. Functional characterization of a novel mutation in NKX2-5 associated with congenital heart disease and adult-onset cardiomyopathy. *Circ Cardiovasc Genet* 2013; **6**: 238-247 [PMID: 23661673 DOI: 10.1161/CIRCGENETICS.113.000057]
- 123 **Qu XK**, Qiu XB, Yuan F, Wang J, Zhao CM, Liu XY, Zhang XL, Li RG, Xu YJ, Hou XM, Fang WY, Liu X, Yang YQ. A novel NKX2.5 loss-of-function mutation associated with congenital bicuspid aortic valve. *Am J Cardiol* 2014; **114**: 1891-1895 [PMID: 25438918 DOI: 10.1016/j.amjcard.2014.09.028]
- 124 **Wang J**, Mao JH, Ding KK, Xu WJ, Liu XY, Qiu XB, Li RG, Qu XK, Xu YJ, Huang RT, Xue S, Yang YQ. A novel NKX2.6 mutation associated with congenital ventricular septal defect. *Pediatr Cardiol* 2015; **36**: 646-656 [PMID: 25380965 DOI: 10.1007/s00246-014-1060-x]
- 125 **Lin Y**, Guo X, Zhao B, Liu J, Da M, Wen Y, Hu Y, Ni B, Zhang K, Yang S, Xu J, Dai J, Wang X, Xia Y, Ma H, Jin G, Yu S, Liu J, Keavney BD, Goodship JA, Cordell HJ, Wang X, Shen H, Sha J, Zhou Z, Chen Y, Mo X, Luo L, Hu Z. Association analysis identifies new risk loci for congenital heart disease in Chinese populations. *Nat Commun* 2015; **6**: 8082 [PMID: 26283027 DOI: 10.1038/ncomms9082]
- 126 **Chowdhury R**, Ashraf H, Melanson M, Tanada Y, Nguyen M, Silberbach M, Wakimoto H, Benson DW, Anderson RH, Kasahara H. Mouse Model of Human Congenital Heart Disease: Progressive Atrioventricular Block Induced by a Heterozygous Nkx2-5 Homeodomain Missense Mutation. *Circ Arrhythm Electrophysiol* 2015; **8**: 1255-1264 [PMID: 26226998]
- 127 **Zhou W**, Zhao L, Jiang JQ, Jiang WF, Yang YQ, Qiu XB. A novel TBX5 loss-of-function mutation associated with sporadic dilated cardiomyopathy. *Int J Mol Med* 2015; **36**: 282-288 [PMID: 25963046 DOI: 10.3892/ijmm.2015.2206]
- 128 **Zhao L**, Ni SH, Liu XY, Wei D, Yuan F, Xu L, Xin-Li RG, Qu XK, Xu YJ, Fang WY, Yang YQ, Qiu XB. Prevalence and spectrum of Nkx2.6 mutations in patients with congenital heart disease. *Eur J Med Genet* 2014; **57**: 579-586 [PMID: 25195019 DOI: 10.1016/j.jmg.2014.08.005]
- 129 **Smith JG**, Magnani JW, Palmer C, Meng YA, Soliman EZ, Musani SK, Kerr KF, Schnabel RB, Lubitz SA, Sotoodehnia N, Redline S, Pfeuffer A, Müller M, Evans DS, Nalls MA, Liu Y, Newman AB, Zonderman AB, Evans MK, Deo R, Ellinor PT, Paltoo DN, Newton-Cheh C, Benjamin EJ, Mehra R, Alonso A, Heckbert SR, Fox ER. Genome-wide association studies of the PR interval in African Americans. *PLoS Genet* 2011; **7**: e1001304 [PMID: 21347284 DOI: 10.1371/journal.pgen.1001304]
- 130 **den Hoed M**, Eijgelsheim M, Esko T, Brundel BJ, Peal DS, Evans DM, Nolte IM, Segrè AV, Holm H, Handsaker RE, Westra HJ, Johnson T, Isaacs A, Yang J, Lundby A, Zhao JH, Kim YJ, Go MJ, Almgren P, Bochud M, Boucher G, Cornelis MC, Gudbjartsson D, Hadley D, van der Harst P, Hayward C, den Heijer M, Igl W, Jackson AU, Kutalik Z, Luan J, Kemp JP, Kristiansson K, Ladenvall C, Lorentzon M, Montasser ME, Njajou OT, O'Reilly PF, Padmanabhan S, St Pourcain B, Rankinen T, Salo P, Tanaka T, Timpson NJ, Vitart V, Waite L, Wheeler W, Zhang W, Draisma HH, Feitosa MF, Kerr KF, Lind PA, Mihailov E, Onland-Moret NC, Song C, Weedon MN, Xie W, Yengo L, Absher D, Albert CM, Alonso A, Arking DE, de Bakker PI, Balkau B, Barlassina C, Benaglio P, Bis JC, Bouatia-Naji N, Brage S, Chanock SJ, Chines PS, Chung M, Darbar D, Dina C, Dörr M, Elliott P, Felix SB, Fischer K, Fuchsberger C, de Geus EJ, Goyette P, Gudnason V, Harris TB, Hartikainen AL, Havulinna AS, Heckbert SR, Hicks AA, Hofman A, Holeywijn S, Hoogstra-Berends F, Hottenga JJ, Jensen MK, Johansson A, Junttila J, Kääb S, Kanon B, Ketkar S, Khaw KT, Knowles JW, Kooner AS, Kors JA, Kumari M, Milani L, Laiho P, Lakatta EG, Langenberg C, Leusink M, Liu Y, Luben RN, Lunetta KL, Lynch SN, Markus MR, Marques-Vidal P, Mateo Leach I, McArdle WL, McCarroll SA, Medland SE, Miller KA, Montgomery GW, Morrison AC, Müller-Nurasyid M, Navarro P, Nelis M, O'Connell JR, O'Donnell CJ, Ong KK, Newman AB, Peters A, Polasek O, Pouta A, Pramstaller PP, Psaty BM, Rao DC, Ring SM, Rossin EJ, Rudan D, Sanna S, Scott RA, Sehmi JS, Sharp S, Shin JT, Singleton AB, Smith AV, Soranzo N, Spector TD, Stewart C, Stringham HM, Tarasov KV, Uitterlinden AG, Vandenput L, Hwang SJ, Whitfield JB, Wijmenga C, Wild SH, Willemssen G, Wilson JF, Witteman JC, Wong A, Wong Q, Jamshidi Y, Zitting P, Boer JM, Boomsma DI, Borecki IB, van Duijn CM, Ekelund U, Forouhi NG, Froguel P, Hingorani A, Ingelsson E, Kivimäki M, Kronmal RA, Kuh D, Lind L, Martin NG, Oostra BA, Pedersen NL, Quertermous T, Rotter JJ, van der Schouw YT, Verschuren WM, Walker M, Albanes D, Arner DO, Assimes TL, Bandinelli S, Boehnke M, de Boer RA, Bouchard C, Caulfield WL, Chambers JC, Curhan G, Cusi D, Eriksson J, Ferrucci L, van Gilst WH, Glorioso N, de Graaf J, Groop L, Gyllenstein U, Hsueh WC, Hu FB, Huikuri HV, Hunter DJ, Iribarren C, Isomaa B, Jarvelin MR, Julia A, Kähönen M, Kiemeny LA, van der Klauw MM, Kooner JS, Kraft P, Iacoviello L, Lehtimäki T, Lokki ML,

- Mitchell BD, Navis G, Nieminen MS, Ohlsson C, Poulter NR, Qi L, Raitakari OT, Rimm EB, Rioux JD, Rizzi F, Rudan I, Salomaa V, Sever PS, Shields DC, Shuldiner AR, Sinisalo J, Stanton AV, Stolk RP, Strachan DP, Tardif JC, Thorsteinsdottir U, Tuomilehto J, van Veldhuisen DJ, Virtamo J, Viikari J, Vollenweider P, Waeber G, Widen E, Cho YS, Olsen JV, Visscher PM, Willer C, Franke L, Erdmann J, Thompson JR, Pfeuffer A, Sotoodehnia N, Newton-Cheh C, Ellinor PT, Stricker BH, Metspalu A, Perola M, Beckmann JS, Smith GD, Stefansson K, Wareham NJ, Munroe PB, Sibon OC, Milan DJ, Snieder H, Samani NJ, Loos RJ. Identification of heart rate-associated loci and their effects on cardiac conduction and rhythm disorders. *Nat Genet* 2013; **45**: 621-631 [PMID: 23583979 DOI: 10.1038/ng.2610]
- 131 **Parvez B**, Shoemaker MB, Muhammad R, Richardson R, Jiang L, Blair MA, Roden DM, Darbar D. Common genetic polymorphism at 4q25 locus predicts atrial fibrillation recurrence after successful cardioversion. *Heart Rhythm* 2013; **10**: 849-855 [PMID: 23428961 DOI: 10.1016/j.hrthm.2013.02.018]
- 132 **Mohanty S**, Santangeli P, Bai R, Di Biase L, Mohanty P, Pump A, Natale A. Variant rs2200733 on chromosome 4q25 confers increased risk of atrial fibrillation: evidence from a meta-analysis. *J Cardiovasc Electrophysiol* 2013; **24**: 155-161 [PMID: 23130627 DOI: 10.1111/jce.12017]
- 133 **Olesen MS**, Holst AG, Jabbari J, Nielsen JB, Christophersen IE, Sajadieh A, Haunsø S, Svendsen JH. Genetic loci on chromosomes 4q25, 7p31, and 12p12 are associated with onset of lone atrial fibrillation before the age of 40 years. *Can J Cardiol* 2012; **28**: 191-195 [PMID: 22336519 DOI: 10.1016/j.cjca.2011.11.016]
- 134 **Henningsen KM**, Olesen MS, Haunsø S, Svendsen JH. Association of rs2200733 at 4q25 with early onset of lone atrial fibrillation in young patients. *Scand Cardiovasc J* 2011; **45**: 324-326 [PMID: 21793630 DOI: 10.3109/14017431.2011.594081]
- 135 **Kilisek M**, Franaszczyk M, Kozluk E, Lodzinski P, Piatkowska A, Broda G, Ploski R, Opolski G. Association between variants on chromosome 4q25, 16q22 and 1q21 and atrial fibrillation in the Polish population. *PLoS One* 2011; **6**: e21790 [PMID: 21760908 DOI: 10.1371/journal.pone.0021790]
- 136 **Chinchilla A**, Daimi H, Lozano-Velasco E, Dominguez JN, Caballero R, Delpón E, Tamargo J, Cinca J, Hove-Madsen L, Aránega AE, Franco D. PITX2 insufficiency leads to atrial electrical and structural remodeling linked to arrhythmogenesis. *Circ Cardiovasc Genet* 2011; **4**: 269-279 [PMID: 21511879 DOI: 10.1161/CIRCGENETICS.110.958116]
- 137 **Sinner MF**, Lubitz SA, Pfeuffer A, Makino S, Beckmann BM, Lunetta KL, Steinbeck G, Perz S, Rahman R, Sonni A, Greenberg SM, Furie KL, Wichmann HE, Meitinger T, Peters A, Benjamin EJ, Rosand J, Ellinor PT, Kääb S. Lack of replication in polymorphisms reported to be associated with atrial fibrillation. *Heart Rhythm* 2011; **8**: 403-409 [PMID: 21056700 DOI: 10.1016/j.hrthm.2010.11.003]
- 138 **Lee KT**, Yeh HY, Tung CP, Chu CS, Cheng KH, Tsai WC, Lu YH, Chang JG, Sheu SH, Lai WT. Association of RS2200733 but not RS10033464 on 4q25 with atrial fibrillation based on the recessive model in a Taiwanese population. *Cardiology* 2010; **116**: 151-156 [PMID: 20606429 DOI: 10.1159/000318172]
- 139 **Wang J**, Klysis E, Sood S, Johnson RL, Wehrens XH, Martin JF. Pitx2 prevents susceptibility to atrial arrhythmias by inhibiting left-sided pacemaker specification. *Proc Natl Acad Sci USA* 2010; **107**: 9753-9758 [PMID: 20457925 DOI: 10.1073/pnas.0912585107]
- 140 **Kirchhof P**, Kahr PC, Kaese S, Piccini I, Vokshi I, Scheld HH, Rotering H, Fortmueller L, Laakmann S, Verheule S, Schotten U, Fabritz L, Brown NA. PITX2c is expressed in the adult left atrium, and reducing Pitx2c expression promotes atrial fibrillation inducibility and complex changes in gene expression. *Circ Cardiovasc Genet* 2011; **4**: 123-133 [PMID: 21282332 DOI: 10.1161/CIRCGENETICS.110.958058]
- 141 **Aguirre LA**, Alonso ME, Badia-Careaga C, Rollán I, Arias C, Fernández-Miñán A, López-Jiménez E, Aránega A, Gómez-Skarmeta JL, Franco D, Manzanares M. Long-range regulatory interactions at the 4q25 atrial fibrillation risk locus involve PITX2c and ENPEP. *BMC Biol* 2015; **13**: 26 [PMID: 25888893 DOI: 10.1186/s12915-015-0138-0]
- 142 **Gore-Panther SR**, Hsu J, Hanna P, Gillinov AM, Pettersson G, Newton DW, Moravec CS, Van Wagoner DR, Chung MK, Barnard J, Smith JD. Atrial Fibrillation associated chromosome 4q25 variants are not associated with PITX2c expression in human adult left atrial appendages. *PLoS One* 2014; **9**: e86245 [PMID: 24465984 DOI: 10.1371/journal.pone.0086245]
- 143 **Martin RI**, Babaei MS, Choy MK, Owens WA, Chico TJ, Keenan D, Yonan N, Koref MS, Keavney BD. Genetic variants associated with risk of atrial fibrillation regulate expression of PITX2, CAV1, MYOZ1, C9orf3 and FANCC. *J Mol Cell Cardiol* 2015; **85**: 207-214 [PMID: 26073630 DOI: 10.1016/j.yjmcc.2015.06.005]
- 144 **Gudbjartsson DF**, Helgason H, Gudjonsson SA, Zink F, Oddson A, Gylfason A, Besenbacher S, Magnusson G, Halldorsson BV, Hjartarson E, Sigurdsson GT, Stacey SN, Frigge ML, Holm H, Saemundsdottir J, Helgadóttir HT, Johannsdóttir H, Sigfusson G, Thorgeirsson G, Sverrisson JT, Gretarsdóttir S, Walters GB, Rafnar T, Thjodleifsson B, Bjornsson ES, Olafsson S, Thorarindottir H, Steingrimsdottir T, Gudmundsdottir TS, Theodors A, Jonasson JG, Sigurdsson A, Bjornsdottir G, Jonsson JJ, Thorarensen O, Ludvigsson P, Gudbjartsson H, Eyjolfsson GI, Sigurdardottir O, Olafsson I, Arnar DO, Magnusson OT, Kong A, Masson G, Thorsteinsdottir U, Helgason A, Sulem P, Stefansson K. Large-scale whole-genome sequencing of the Icelandic population. *Nat Genet* 2015; **47**: 435-444 [PMID: 25807286 DOI: 10.1038/ng.3247]
- 145 **Li C**, Wang F, Yang Y, Fu F, Xu C, Shi L, Li S, Xia Y, Wu G, Cheng X, Liu H, Wang C, Wang P, Hao J, Ke Y, Zhao Y, Liu M, Zhang R, Gao L, Yu B, Zeng Q, Liao Y, Yang B, Tu X, Wang QK. Significant association of SNP rs2106261 in the ZFX3 gene with atrial fibrillation in a Chinese Han GenElD population. *Hum Genet* 2011; **129**: 239-246 [PMID: 21107608 DOI: 10.1007/s00439-010-0912-6]
- 146 **Zhai C**, Cong H, Liu Y, Zhang Y, Liu X, Zhang H, Ren Z. Rs7193343 polymorphism in zinc finger homeobox 3 (ZFX3) gene and atrial fibrillation: an updated meta-analysis of 10 case-control comparisons. *BMC Cardiovasc Disord* 2015; **15**: 58 [PMID: 26112950 DOI: 10.1186/s12872-015-0044-y]
- 147 **Lin H**, Sinner MF, Brody JA, Arking DE, Lunetta KL, Rienstra M, Lubitz SA, Magnani JW, Sotoodehnia N, McKnight B, McManus DD, Boerwinkle E, Psaty BM, Rotter JJ, Bis JC, Gibbs RA, Muzny D, Kovar CL, Morrison AC, Gupta M, Folsom AR, Kääb S, Heckbert SR, Alonso A, Ellinor PT, Benjamin EJ. Targeted sequencing in candidate genes for atrial fibrillation: the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Targeted Sequencing Study. *Heart Rhythm* 2014; **11**: 452-457 [PMID: 24239840 DOI: 10.1016/j.hrthm.2013.11.012]
- 148 **Mahida S**, Mills RW, Tucker NR, Simonson B, Macri V, Lemoine MD, Das S, Milan DJ, Ellinor PT. Overexpression of KCNN3 results in sudden cardiac death. *Cardiovasc Res* 2014; **101**: 326-334 [PMID: 24296650 DOI: 10.1093/cvr/cvt269]
- 149 **Lozano-Velasco E**, Hernández-Torres F, Daimi H, Serra SA, Herraiz A, Hove-Madsen L, Aránega A, Franco D. Pitx2 impairs calcium handling in a dose-dependent manner by modulating Wnt signalling. *Cardiovasc Res* 2016; **109**: 55-66 [PMID: 26243430]
- 150 **Jiang Q**, Ni B, Shi J, Han Z, Qi R, Xu W, Wang D, Wang DW, Chen M. Down-regulation of ATBF1 activates STAT3 signaling via PIAS3 in pacing-induced HL-1 atrial myocytes. *Biochem Biophys Res Commun* 2014; **449**: 278-283 [PMID: 24857987 DOI: 10.1016/j.bbrc.2014.05.041]
- 151 **Kääb S**, Darbar D, van Noord C, Dupuis J, Pfeuffer A, Newton-Cheh C, Schnabel R, Makino S, Sinner MF, Kannankeril PJ, Beckmann BM, Choudry S, Donahue BS, Heeringa J, Perz S, Lunetta KL, Larson MG, Levy D, MacRae CA, Ruskin JN, Wacker A, Schömig A, Wichmann HE, Steinbeck G, Meitinger T, Uitterlinden AG, Witteman JC, Roden DM, Benjamin EJ, Ellinor PT. Large scale replication and meta-analysis of variants on chromosome 4q25 associated with atrial fibrillation. *Eur Heart J* 2009; **30**: 813-819 [PMID: 19141561 DOI: 10.1093/eurheartj/

ehn578]

- 152 **Tsai CT**, Hsieh CS, Chang SN, Chuang EY, Juang JM, Lin LY, Lai LP, Hwang JJ, Chiang FT, Lin JL. Next-generation sequencing of nine atrial fibrillation candidate genes identified novel de novo mutations in patients with extreme trait of atrial fibrillation. *J Med Genet* 2015; **52**: 28-36 [PMID: 25391453 DOI: 10.1136/jmedgenet-2014-102618]
- 153 **Huang Y**, Wang C, Yao Y, Zuo X, Chen S, Xu C, Zhang H, Lu Q, Chang L, Wang F, Wang P, Zhang R, Hu Z, Song Q, Yang X, Li C, Li S, Zhao Y, Yang Q, Yin D, Wang X, Si W, Li X, Xiong X, Wang D, Huang Y, Luo C, Li J, Wang J, Chen J, Wang L, Wang L, Han M, Ye J, Chen F, Liu J, Liu Y, Wu G, Yang B, Cheng X, Liao Y, Wu Y, Ke T, Chen Q, Tu X, Elston R, Rao S, Yang Y, Xia Y, Wang QK. Molecular Basis of Gene-Gene Interaction: Cyclic Cross-Regulation of Gene Expression and Post-GWAS Gene-Gene Interaction Involved in Atrial Fibrillation. *PLoS Genet* 2015; **11**: e1005393 [PMID: 26267381 DOI: 10.1371/journal.pgen.1005393]
- 154 **Li G**, Zhang R, Gao L, Zhang S, Dong Y, Yin X, Chang D, Yang Y, Xia Y. Lack of association between rs3807989 in *cav1* and atrial fibrillation. *Int J Clin Exp Pathol* 2014; **7**: 4339-4344 [PMID: 25120818]
- 155 **Tada H**, Shiffman D, Smith JG, Sjögren M, Lubitz SA, Ellinor PT, Louie JZ, Catanese JJ, Engström G, Devlin JJ, Kathiresan S, Melander O. Twelve-single nucleotide polymorphism genetic risk score identifies individuals at increased risk for future atrial fibrillation and stroke. *Stroke* 2014; **45**: 2856-2862 [PMID: 25123217 DOI: 10.1161/STROKEAHA.114.006072]
- 156 **Pfeufer A**, van Noord C, Marciante KD, Arking DE, Larson MG, Smith AV, Tarasov KV, Müller M, Sotoodehnia N, Sinner MF, Verwoert GC, Li M, Kao WH, Köttgen A, Coresh J, Bis JC, Psaty BM, Rice K, Rotter JJ, Rivadeneira F, Hofman A, Kors JA, Stricker BH, Uitterlinden AG, van Duijn CM, Beckmann BM, Sauter W, Gieger C, Lubitz SA, Newton-Cheh C, Wang TJ, Magnani JW, Schnabel RB, Chung MK, Barnard J, Smith JD, Van Wagoner DR, Vasan RS, Aspelund T, Eiriksdottir G, Harris TB, Launer LJ, Najjar SS, Lakatta E, Schlessinger D, Uda M, Abecasis GR, Müller-Myhok B, Ehret GB, Boerwinkle E, Chakravarti A, Soliman EZ, Lunetta KL, Perz S, Wichmann HE, Meitinger T, Levy D, Gudnason V, Ellinor PT, Sanna S, Kääb S, Witteman JC, Alonso A, Benjamin EJ, Heckbert SR. Genome-wide association study of PR interval. *Nat Genet* 2010; **42**: 153-159 [PMID: 20062060 DOI: 10.1038/ng.517]
- 157 **Holm H**, Gudbjartsson DF, Arnar DO, Thorleifsson G, Thorgeirsson G, Stefansson H, Gudjonsson SA, Jonasdottir A, Mathiesen EB, Njølstad I, Nyrnes A, Wilsgaard T, Hald EM, Hveem K, Stoltenberg C, Løchen ML, Kong A, Thorsteinsdottir U, Stefansson K. Several common variants modulate heart rate, PR interval and QRS duration. *Nat Genet* 2010; **42**: 117-122 [PMID: 20062063 DOI: 10.1038/ng.511]
- 158 **Butler AM**, Yin X, Evans DS, Nalls MA, Smith EN, Tanaka T, Li G, Buxbaum SG, Whitsel EA, Alonso A, Arking DE, Benjamin EJ, Berenson GS, Bis JC, Chen W, Deo R, Ellinor PT, Heckbert SR, Heiss G, Hsueh WC, Keating BJ, Kerr KF, Li Y, Limacher MC, Liu Y, Lubitz SA, Marciante KD, Mehra R, Meng YA, Newman AB, Newton-Cheh C, North KE, Palmer CD, Psaty BM, Quibrera PM, Redline S, Reiner AP, Rotter JJ, Schnabel RB, Schork NJ, Singleton AB, Smith JG, Soliman EZ, Srinivasan SR, Zhang ZM, Zonderman AB, Ferrucci L, Murray SS, Evans MK, Sotoodehnia N, Magnani JW, Avery CL. Novel loci associated with PR interval in a genome-wide association study of 10 African American cohorts. *Circ Cardiovasc Genet* 2012; **5**: 639-646 [PMID: 23139255 DOI: 10.1161/CIRCGENETICS.112.963991]
- 159 **Bezzina CR**, Barc J, Mizusawa Y, Remme CA, Gourraud JB, Simonet F, Verkerk AO, Schwartz PJ, Crotti L, Dagradi F, Guicheney P, Fressart V, Leenhardt A, Antzelevitch C, Bartkowiak S, Borggreffe M, Schimpf R, Schulze-Bahr E, Zumhagen S, Behr ER, Bastiaenen R, Tfelt-Hansen J, Olesen MS, Kääb S, Beckmann BM, Weeke P, Watanabe H, Endo N, Minamino T, Horie M, Ohno S, Hasegawa K, Makita N, Nogami A, Shimizu W, Aiba T, Froguel P, Balkau B, Lantieri O, Torchio M, Wiese C, Weber D, Wolswinkel R, Coronel R, Boukens BJ, Béziau S, Charpentier E, Chatel S, Despres A, Gros F, Kyndt F, Lecoite S, Lindenbaum P, Portero V, Violleau J, Gessler M, Tan HL, Roden DM, Christoffels VM, Le Marec H, Wilde AA, Probst V, Schott JJ, Dina C, Redon R. Common variants at SCN5A-SCN10A and HEY2 are associated with Brugada syndrome, a rare disease with high risk of sudden cardiac death. *Nat Genet* 2013; **45**: 1044-1049 [PMID: 23872634 DOI: 10.1038/ng.2712]
- 160 **Newton-Cheh C**, Eijgelsheim M, Rice KM, de Bakker PI, Yin X, Estrada K, Bis JC, Marciante K, Rivadeneira F, Noseworthy PA, Sotoodehnia N, Smith NL, Rotter JJ, Kors JA, Witteman JC, Hofman A, Heckbert SR, O'Donnell CJ, Uitterlinden AG, Psaty BM, Lumley T, Larson MG, Stricker BH. Common variants at ten loci influence QT interval duration in the QTGEN Study. *Nat Genet* 2009; **41**: 399-406 [PMID: 19305408 DOI: 10.1038/ng.364]
- 161 **Pfeufer A**, Sanna S, Arking DE, Müller M, Gateva V, Fuchsberger C, Ehret GB, Orrú M, Pattaro C, Köttgen A, Perz S, Usala G, Barbalic M, Li M, Pütz B, Scuteri A, Prineas RJ, Sinner MF, Gieger C, Najjar SS, Kao WH, Mühleisen TW, Dei M, Happle C, Möhlenkamp S, Crisponi L, Erbel R, Jöckel KH, Naitza S, Steinbeck G, Marroni F, Hicks AA, Lakatta E, Müller-Myhok B, Pramstaller PP, Wichmann HE, Schlessinger D, Boerwinkle E, Meitinger T, Uda M, Coresh J, Kääb S, Abecasis GR, Chakravarti A. Common variants at ten loci modulate the QT interval duration in the QTSCD Study. *Nat Genet* 2009; **41**: 407-414 [PMID: 19305409 DOI: 10.1038/ng.362]
- 162 **Arking DE**, Pulit SL, Crotti L, van der Harst P, Munroe PB, Koopmann TT, Sotoodehnia N, Rossin EJ, Morley M, Wang X, Johnson AD, Lundby A, Gudbjartsson DF, Noseworthy PA, Eijgelsheim M, Bradford Y, Tarasov KV, Dörr M, Müller-Nurasyid M, Lahtinen AM, Nolte IM, Smith AV, Bis JC, Isaacs A, Newhouse SJ, Evans DS, Post WS, Waggott D, Lyytikäinen LP, Hicks AA, Eisele L, Ellinghaus D, Hayward C, Navarro P, Ulivi S, Tanaka T, Tester DJ, Chatel S, Gustafsson S, Kumari M, Morris RW, Naluai ÁT, Padmanabhan S, Kluttig A, Strohmer B, Panayiotou AG, Torres M, Knoflach M, Hubacek JA, Slowikowski K, Raychaudhuri S, Kumar RD, Harris TB, Launer LJ, Shuldiner AR, Alonso A, Bader JS, Ehret G, Huang H, Kao WH, Strait JB, Macfarlane PW, Brown M, Caulfield MJ, Samani NJ, Kronenberg F, Willeit J, Smith JG, Greiser KH, Meyer Zu Schwabedissen H, Werdan K, Carella M, Zelante L, Heckbert SR, Psaty BM, Rotter JJ, Kolcic I, Polašek O, Wright AF, Griffin M, Daly MJ, Arnar DO, Hólm H, Thorsteinsdottir U, Denny JC, Roden DM, Zuvich RL, Emilsson V, Plump AS, Larson MG, O'Donnell CJ, Yin X, Bobbo M, D'Adamo AP, Iorio A, Sinagra G, Carracedo A, Cummings SR, Nalls MA, Jula A, Kontula KK, Marjamaa A, Oikarinen L, Perola M, Porthan K, Erbel R, Hoffmann P, Jöckel KH, Kälisch H, Nöthen MM, den Hoed M, Loos RJ, Thelle DS, Gieger C, Meitinger T, Perz S, Peters A, Prucha R, Sinner MF, Waldenberger M, de Boer RA, Franke L, van der Vleuten P, Beckmann BM, Martens E, Bardai A, Hofman N, Wilde AA, Behr ER, Dalageorgou C, Giudicessi JR, Medeiros-Domingo A, Barc J, Kyndt F, Probst V, Ghidoni A, Insolia R, Hamilton RM, Scherer SW, Brandimarto J, Margulies K, Moravec CE, del Greco M F, Fuchsberger C, O'Connell JR, Lee WK, Watt GC, Campbell H, Wild SH, El Mokhtari NE, Frey N, Asselbergs FW, Mateo Leach I, Navis G, van den Berg MP, van Veldhuisen DJ, Kellis M, Krijthe BP, Franco OH, Hofman A, Kors JA, Uitterlinden AG, Witteman JC, Kedenko L, Lamina C, Oostra BA, Abecasis GR, Lakatta EG, Mulas A, Orrú M, Schlessinger D, Uda M, Markus MR, Völker U, Snieder H, Spector TD, Ärnlöv J, Lind L, Sundström J, Syvänen AC, Kivimäki M, Kähönen M, Mononen N, Raitakari OT, Viikari JS, Adamkova V, Kiechl S, Brion M, Nicolaides AN, Paulweber B, Haerting J, Dominiczak AF, Nyberg F, Whincup PH, Hingorani AD, Schott JJ, Bezzina CR, Ingelsson E, Ferrucci L, Gasparini P, Wilson JF, Rudan I, Franke A, Mühleisen TW, Pramstaller PP, Lehtimäki TJ, Paterson AD, Parsa A, Liu Y, van Duijn CM, Siscovick DS, Gudnason V, Jamshidi Y, Salomaa V, Felix SB, Sanna S, Ritchie MD, Stricker BH, Stefansson K, Boyer LA, Cappola TP, Olsen JV, Lage K, Schwartz PJ, Kääb S, Chakravarti A, Ackerman MJ, Pfeufer A, de Bakker PI, Newton-Cheh C. Genetic association

- study of QT interval highlights role for calcium signaling pathways in myocardial repolarization. *Nat Genet* 2014; **46**: 826-836 [PMID: 24952745 DOI: 10.1038/ng.3014]
- 163 **Li L**, Shen C, Yao Z, Liang J, Huang C. Genetic variants of potassium voltage-gated channel genes (KCNQ1, KCNH2, and KCNE1) affected the risk of atrial fibrillation in elderly patients. *Genet Test Mol Biomarkers* 2015; **19**: 359-365 [PMID: 26066992 DOI: 10.1089/gtmb.2014.0307]
- 164 **Chu HM**, Feng MJ, Li YG, Zhang YX, Ma JF, He B, Yu YB, Liu J, Chen XM. Polymorphisms but not mutations of the KCNQ1 gene are associated with lone atrial fibrillation in the Chinese Han population. *ScientificWorldJournal* 2013; **2013**: 373454 [PMID: 23710137 DOI: 10.1155/2013/373454]
- 165 **Zeng Z**, Tan C, Teng S, Chen J, Su S, Zhou X, Wang F, Zhang S, Gu D, Makielski JC, Pu J. The single nucleotide polymorphisms of I(Ks) potassium channel genes and their association with atrial fibrillation in a Chinese population. *Cardiology* 2007; **108**: 97-103 [PMID: 17016049]
- 166 **Jabbari J**, Olesen MS, Holst AG, Nielsen JB, Haunso S, Svendsen JH. Common polymorphisms in KCNJ5 [corrected] are associated with early-onset lone atrial fibrillation in Caucasians. *Cardiology* 2011; **118**: 116-120 [PMID: 21555883 DOI: 10.1159/000323840]
- 167 **Hauer RN**, Groenewegen WA, Firouzi M, Ramanna H, Jongsma HJ. Cx40 polymorphism in human atrial fibrillation. *Adv Cardiol* 2006; **42**: 284-291 [PMID: 16646598]
- 168 **Zang X**, Zhang S, Xia Y, Li S, Fu X, Wang F, Zhang R, Tian X, Gao L, Zhang J, Yang Y, Tu X, Wang Q. SNP rs3825214 in TBX5 is associated with lone atrial fibrillation in Chinese Han population. *PLoS One* 2013; **8**: e64966 [PMID: 23717681 DOI: 10.1371/journal.pone.0064966]
- 169 **Wang C**, Wu M, Qian J, Li B, Tu X, Xu C, Li S, Chen S, Zhao Y, Huang Y, Shi L, Cheng X, Liao Y, Chen Q, Xia Y, Yao W, Wu G, Cheng M, Wang QK. Identification of rare variants in TNNI3 with atrial fibrillation in a Chinese GenEd population. *Mol Genet Genomics* 2015; **291**: 79-92 [PMID: 26169204]
- 170 **Jabbari J**, Olesen MS, Yuan L, Nielsen JB, Liang B, Macri V, Christophersen IE, Nielsen N, Sajadieh A, Ellinor PT, Grunnet M, Haunso S, Holst AG, Svendsen JH, Jespersen T. Common and rare variants in SCN10A modulate the risk of atrial fibrillation. *Circ Cardiovasc Genet* 2015; **8**: 64-73 [PMID: 25691686 DOI: 10.1161/HCG.0000000000000022]
- 171 **Delaney JT**, Muhammad R, Shi Y, Schilderout JS, Blair M, Short L, Roden DM, Darbar D. Common SCN10A variants modulate PR interval and heart rate response during atrial fibrillation. *Europace* 2014; **16**: 485-490 [PMID: 24072447 DOI: 10.1093/europace/eut278]
- 172 **Franco D**, Aranega A. Post-transcriptional regulatory mechanisms. Clinic, Genetics and Molecular Pathways of Congenital Heart Diseases. In: Sperling S, Driscoll D, Kelly R, editors. Springer, 2015; In press
- 173 **Espinoza-Lewis RA**, Wang DZ. MicroRNAs in heart development. *Curr Top Dev Biol* 2012; **100**: 279-317 [PMID: 22449848 DOI: 10.1016/B978-0-12-387786-4.00009-9]
- 174 **Zhou J**, Dong X, Zhou Q, Wang H, Qian Y, Tian W, Ma D, Li X. microRNA expression profiling of heart tissue during fetal development. *Int J Mol Med* 2014; **33**: 1250-1260 [PMID: 24604530 DOI: 10.3892/ijmm.2014.1691]
- 175 **Hu DL**, Liu YQ, Chen FK, Sheng YH, Yang R, Kong XQ, Cao KJ, Zhang JS, Qian LM. Differential expression of microRNAs in cardiac myocytes compared to undifferentiated P19 cells. *Int J Mol Med* 2011; **28**: 59-64 [PMID: 21455566 DOI: 10.3892/ijmm.2011.664]
- 176 **Synergren J**, Améen C, Lindahl A, Olsson B, Sartipy P. Expression of microRNAs and their target mRNAs in human stem cell-derived cardiomyocyte clusters and in heart tissue. *Physiol Genomics* 2011; **43**: 581-594 [PMID: 20841501 DOI: 10.1152/physiolgenomics.00074.2010]
- 177 **Nishi H**, Sakaguchi T, Miyagawa S, Yoshikawa Y, Fukushima S, Saito S, Ueno T, Kuratani T, Sawa Y. Impact of microRNA expression in human atrial tissue in patients with atrial fibrillation undergoing cardiac surgery. *PLoS One* 2013; **8**: e73397 [PMID: 24069193 DOI: 10.1371/journal.pone.0073397]
- 178 **Wang J**, Wang Y, Han J, Li Y, Xie C, Xie L, Shi J, Zhang J, Yang B, Chen D, Meng X. Integrated analysis of microRNA and mRNA expression profiles in the left atrium of patients with nonvalvular paroxysmal atrial fibrillation: Role of miR-146b-5p in atrial fibrosis. *Heart Rhythm* 2015; **12**: 1018-1026 [PMID: 25617731 DOI: 10.1016/j.hrthm.2015.01.026]
- 179 **Li M**, Zhang J. Circulating MicroRNAs: Potential and Emerging Biomarkers for Diagnosis of Cardiovascular and Cerebrovascular Diseases. *Biomed Res Int* 2015; **2015**: 730535 [PMID: 26180810 DOI: 10.1155/2015/730535]
- 180 **Orenes-Piñero E**, Montoro-García S, Patel JV, Valdés M, Marín F, Lip GY. Role of microRNAs in cardiac remodelling: new insights and future perspectives. *Int J Cardiol* 2013; **167**: 1651-1659 [PMID: 23063140 DOI: 10.1016/j.ijcard.2012.09.120]
- 181 **Poudel P**, Xu Y, Cui Z, Sharma D, Tian B, Paudel S. Atrial fibrillation: recent advances in understanding the role of microRNAs in atrial remodeling with an electrophysiological overview. *Cardiology* 2015; **131**: 58-67 [PMID: 25871909 DOI: 10.1159/000375403]
- 182 **Gomes da Silva AM**, Silbiger VN. miRNAs as biomarkers of atrial fibrillation. *Biomarkers* 2014; **19**: 631-636 [PMID: 25171770 DOI: 10.3109/1354750X.2014.954001]
- 183 **Liu Z**, Zhou C, Liu Y, Wang S, Ye P, Miao X, Xia J. The expression levels of plasma microRNAs in atrial fibrillation patients. *PLoS One* 2012; **7**: e44906 [PMID: 23028671 DOI: 10.1371/journal.pone.0044906]
- 184 **van den Boogaard M**, Smemo S, Burnicka-Turek O, Arnolds DE, van de Werken HJ, Klous P, McKean D, Muehlschlegel JD, Moosmann J, Toka O, Yang XH, Koopmann TT, Adriaens ME, Bezzina CR, de Laat W, Seidman C, Seidman JG, Christoffels VM, Nobrega MA, Barnett P, Moskowitz IP. A common genetic variant within SCN10A modulates cardiac SCN5A expression. *J Clin Invest* 2014; **124**: 1844-1852 [PMID: 24642470 DOI: 10.1172/JCI73140]
- 185 **Tao Y**, Zhang M, Li L, Bai Y, Zhou Y, Moon AM, Kaminski HJ, Martin JF. Pitx2, an atrial fibrillation predisposition gene, directly regulates ion transport and intercalated disc genes. *Circ Cardiovasc Genet* 2014; **7**: 23-32 [PMID: 24395921 DOI: 10.1161/CIRCGENETICS.113.000259]
- 186 **Nagy II**, Railo A, Rapila R, Hast T, Sormunen R, Tavi P, Räsänen J, Vainio SJ. Wnt-11 signalling controls ventricular myocardium development by patterning N-cadherin and beta-catenin expression. *Cardiovasc Res* 2010; **85**: 100-109 [PMID: 19622544 DOI: 10.1093/cvr/cvp254]
- 187 **Pandur P**, Läsche M, Eisenberg LM, Kühl M. Wnt-11 activation of a non-canonical Wnt signalling pathway is required for cardiogenesis. *Nature* 2002; **418**: 636-641 [PMID: 12167861]
- 188 **Martin A**, Maher S, Summerhust K, Davidson D, Murphy P. Differential deployment of paralogous Wnt genes in the mouse and chick embryo during development. *Evol Dev* 2012; **14**: 178-195 [PMID: 23017026 DOI: 10.1111/j.1525-142X.2012.00534.x]
- 189 **Wang J**, Bai Y, Li N, Ye W, Zhang M, Greene SB, Tao Y, Chen Y, Wehrens XH, Martin JF. Pitx2-microRNA pathway that delimits sinoatrial node development and inhibits predisposition to atrial fibrillation. *Proc Natl Acad Sci USA* 2014; **111**: 9181-9186 [PMID: 24927531 DOI: 10.1073/pnas.1405411111]
- 190 **Chiang DY**, Kongchan N, Beavers DL, Alsina KM, Voigt N, Neilson JR, Jakob H, Martin JF, Dobrev D, Wehrens XH, Li N. Loss of microRNA-106b-25 cluster promotes atrial fibrillation by enhancing ryanodine receptor type-2 expression and calcium release. *Circ Arrhythm Electrophysiol* 2014; **7**: 1214-1222 [PMID: 25389315 DOI: 10.1161/CIRCEP.114.001973]
- 191 **Barana A**, Matamoros M, Dolz-Gaitón P, Pérez-Hernández M, Amorós I, Núñez M, Sacristán S, Pedraz Á, Pinto Á, Fernández-Avilés F, Tamargo J, Delpón E, Caballero R. Chronic atrial fibrillation increases microRNA-21 in human atrial myocytes decreasing L-type calcium current. *Circ Arrhythm Electrophysiol* 2014; **7**: 861-868 [PMID: 25107449 DOI: 10.1161/CIRCEP.114.001709]
- 192 **Daimi H**, Lozano-Velasco E, Haj Khelil A, Chibani JB, Barana A, Amorós I, González de la Fuente M, Caballero R, Aranega

- A, Franco D. Regulation of SCN5A by microRNAs: miR-219 modulates SCN5A transcript expression and the effects of flecainide intoxication in mice. *Heart Rhythm* 2015; **12**: 1333-1342 [PMID: 25701775 DOI: 10.1016/j.hrthm.2015.02.018]
- 193 **Luo X**, Pan Z, Shan H, Xiao J, Sun X, Wang N, Lin H, Xiao L, Maguy A, Qi XY, Li Y, Gao X, Dong D, Zhang Y, Bai Y, Ai J, Sun L, Lu H, Luo XY, Wang Z, Lu Y, Yang B, Nattel S. MicroRNA-26 governs profibrillatory inward-rectifier potassium current changes in atrial fibrillation. *J Clin Invest* 2013; **123**: 1939-1951 [PMID: 23543060 DOI: 10.1172/JCI62185]
- 194 **Jia X**, Zheng S, Xie X, Zhang Y, Wang W, Wang Z, Zhang Y, Wang J, Gao M, Hou Y. MicroRNA-1 accelerates the shortening of atrial effective refractory period by regulating KCNE1 and KCNB2 expression: an atrial tachypacing rabbit model. *PLoS One* 2013; **8**: e85639 [PMID: 24386485 DOI: 10.1371/journal.pone.0085639]
- 195 **Li YD**, Hong YF, Yusufuaji Y, Tang BP, Zhou XH, Xu GJ, Li JX, Sun L, Zhang JH, Xin Q, Xiong J, Ji YT, Zhang Y. Altered expression of hyperpolarization-activated cyclic nucleotide-gated channels and microRNA-1 and -133 in patients with age-associated atrial fibrillation. *Mol Med Rep* 2015; **12**: 3243-3248 [PMID: 26005035 DOI: 10.3892/mmr.2015.3831]
- 196 **Mommersteeg MT**, Brown NA, Prall OW, de Gier-de Vries C, Harvey RP, Moorman AF, Christoffels VM. Pitx2c and Nkx2-5 are required for the formation and identity of the pulmonary myocardium. *Circ Res* 2007; **101**: 902-909 [PMID: 17823370]
- 197 **Mints Y**, Yarmohammadi H, Khurram IM, Hoyt H, Hansford R, Zimmerman SL, Steinberg SJ, Judge DP, Tomaselli GF, Calkins H, Zipunnikov V, Nazarian S. Association of common variations on chromosome 4q25 and left atrial volume in patients with atrial fibrillation. *Clin Med Insights Cardiol* 2015; **9**: 39-45 [PMID: 26005361 DOI: 10.4137/CMC.S21712]
- 198 **Scridon A**, Fouilloux-Meugnier E, Loizon E, Rome S, Julien C, Barrès C, Chevalier P. Long-standing arterial hypertension is associated with Pitx2 down-regulation in a rat model of spontaneous atrial tachyarrhythmias. *Europace* 2015; **17**: 160-165 [PMID: 24908044 DOI: 10.1093/europace/euu139]
- 199 **Torrado M**, Franco D, Hernández-Torres F, Crespo-Leiro MG, Iglesias-Gil C, Castro-Beiras A, Mikhailov AT. Pitx2c is reactivated in the failing myocardium and stimulates myf5 expression in cultured cardiomyocytes. *PLoS One* 2014; **9**: e90561 [PMID: 24595098 DOI: 10.1371/journal.pone.0090561]
- 200 **Torrado M**, Franco D, Lozano-Velasco E, Hernández-Torres F, Calviño R, Aldama G, Centeno A, Castro-Beiras A, Mikhailov A. A MicroRNA-Transcription Factor Blueprint for Early Atrial Arrhythmogenic Remodeling. *Biomed Res Int* 2015; **2015**: 263151 [PMID: 26221584 DOI: 10.1155/2015/263151]

P- Reviewer: Kettering K, Liu T, Said SAM **S- Editor:** Ji FF

L- Editor: A **E- Editor:** Lu YJ





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>



World Journal of *Medical Genetics*

World J Med Genet 2016 May 27; 6(2): 17-21





Editorial Board

2016-2019

The *World Journal of Medical Genetics* Editorial Board consists of 331 members, representing a team of worldwide experts in medical genetics. They are from 50 countries, including Argentina (1), Armenia (1), Australia (4), Austria (2), Belgium (2), Brazil (6), Bulgaria (2), Canada (6), Chile (1), China (39), Croatia (1), Czech Republic (1), Denmark (3), Ecuador (1), Egypt (2), France (10), Germany (4), Greece (2), Hungary (1), India (13), Iran (7), Ireland (1), Israel (2), Italy (22), Japan (23), Malaysia (2), Mexico (2), Netherlands (5), New Zealand (2), Norway (2), Pakistan (1), Peru (1), Poland (1), Portugal (1), Russia (2), Saudi Arabia (4), Serbia (1), Singapore (3), Slovenia (2), South Korea (7), Spain (16), Sweden (1), Switzerland (2), Thailand (3), Turkey (8), Ukraine (1), United Arab Emirates (1), United Kingdom (10), United States (95) and Uruguay (1).

EDITOR-IN-CHIEF

Hans van Bokhoven, *Nijmegen*

GUEST EDITORIAL BOARD MEMBERS

Chia-Hsiang Chen, *Zhunan*
Ji-Yih Chen, *Keelung*
Bor-Luen Chiang, *Taipei*
Yu-Chen Hu, *Hsinchu*
Suh-Hang H Juo, *Kaohsiung*
Hsien-Hsiung Lee, *Taichung*
Yueh-Lun Lee, *Taipei*
Chin-San Liu, *Changhua*
Pei-Jung Lu, *Tainan*
Chien-Kuo Tai, *Chiayi*
Po-Nien Tsao, *Taipei*
Wendy Wen-Hui Weng, *Taipei*
Kuo-Liang Yang, *Hualien*

MEMBERS OF THE EDITORIAL BOARD



Argentina

Carlos J Pirola, *Buenos Aires*



Armenia

Hasmik Mkrtchyan, *Yerevan*



Australia

Simon Easteal, *Canberra*
Jeremy BM Jowett, *Melbourne*

Karen Anne Mather, *Randwick*
Grant Morahan, *Perth*



Austria

Harald N Aschauer, *Vienna*
Anton Wutz, *Vienna*



Belgium

Teresinha Leal, *Brussels*
van West, Dirk, *Antwerp*



Brazil

Pereira Priscila Krauss, *Rio de Janeiro*
Sharbel Weidner Maluf, *Porto Alegre*
Milton Ozório Moraes, *Rio de Janeiro*
Fernando Regla Vargas, *Rio de Janeiro*
Jeane Eliete Laguila Visentainer, *Sao Paulo*
Isabela Jube Wastowski, *Goiania*



Bulgaria

Spaska Angelova Stanilova, *Stara Zagora*
Albena Todorova, *Sofia*



Canada

Régen Drouin, *Sherbrooke*
Gonzalo Hortelano, *Hamilton*
William Jia, *Vancouver*

Sean Li, *Ottawa*
Shiva M Singh, *London*
Liting Song, *Toronto*



Chile

Jara Sosa Lilian Elena, *Santiago*



China

George G Chen, *Hong Kong*
Ning-Han Feng, *Nanjing*
Shi-Ang Huang, *Wuhan*
Wei Huang, *Shanghai*
Chun-Yan Ji, *Jinan*
Hong-Chuan Jin, *Hangzhou*
Ke-Shen Li, *Zhanjiang*
Zhuo-Zhuang Lu, *Beijing*
Xiao-Ping Luo, *Wuhan*
Wen-Bin Qian, *Hangzhou*
Rong Rui, *Nanjing*
Lun-Quan Sun, *Changsha*
Qian Tao, *Hong Kong*
George Sai-Wah Tsao, *Hong Kong*
Ming-Rong Wang, *Beijing*
Shi-Jie Wang, *Shijiazhuang*
Wei Wang, *Beijing*
Ji Wu, *Shanghai*
Hua-Xi Xu, *Zhenjiang*
Ruian Xu, *Xiamen*
Wei-Hua Yan, *Linhai*
Rong-Cun Yang, *Tianjin*
Bang-Ce Ye, *Shanghai*
Rui-Xing Yin, *Nanning*
Yi Zeng, *Beijing*
Xiao-Feng Zhu, *Guangzhou*

**Croatia**

Gordan Lauc, *Zagreb*

**Czech Republic**

Marie Lipoldová, *Prague*

**Denmark**

Vibeke Andersen, *Odense*
Thomas V. O. Hansen, *Copenhagen*
Goutam Sahana, *Tjele*

**Ecuador**

Cesar Paz-Y-Mino, *Quito*

**Egypt**

Menha MA Swellam, *Giza*
Samia Ali Temtamy, *Cairo*

**France**

Christophe Chevillard, *Marseille*
Johanna Chluba, *Dijon*
Zdenko Herceg, *Lyon*
Enzo Lalli, *Valbonne*
James Lespinasse, *Chambéry*
Bernard S. Lopez, *Fontenay aux Roses*
Eric Pasmant, *Clichy*
Georges Vassaux, *Nice Cedex*
Kay-Dietrich Wagner, *Nice Cedex*
Qing Wang, *Lyon*

**Germany**

Stefan Bohringer, *Essen*
Anibh Martin Das, *Hannover*
Gudrun A. Rappold, *Heidelberg*
Hermona Soreq, *Jerusalem*

**Greece**

Roxani Angelopoulou, *Athens*
Voula Velissariou, *Athens*

**Hungary**

Béla Meleg, *Pecs*

**India**

Arvind Kumar Arya, *Meerut*
Prakash Sadashiv Gambhir, *Pune*
Katta KM Girisha, *Karnataka*
KAISER Jamil, *Hyderabad*

Narendra Joshi, *Navi Mumbai*
Swarna Mandava, *Mumbai*
Chinmay K Panda, *Kolkata*
Rajni Rani, *New Delhi*
Susanta Roychoudhury, *Kolkata*
Sudeep R Shah, *Mumbai*
Ranbir Chander Sobti, *Chandigarh*
Meera Vaswani, *New Delhi*
Suresh P Vyas, *Sagar*

**Iran**

Yahya Daneshbod, *Shiraz*
Fariborz Ghaffarpasand, *Shiraz*
Mohammad H Modarressi, *Tehran*
Shirin Hasani Ranjbar, *Tehran*
Nader tajik, *Tehran*
Dor Mohammad Kordi Tamandani, *Zahedan*
Alireza Zomorodipour, *Tehra*

**Ireland**

Ross McManus, *Dublin*

**Israel**

Aliza Shlomit Amiel, *Kfar-Saba*
Rachel Ben-Shlomo, *Tivon*

**Italy**

Francesco Acquati, *Varese*
Nenad Bukvic, *Bari*
Gabriele Candiani, *Milan*
Antonio Cao, *Cagliari*
Raffaella Cascella, *Rome*
Marco Castori, *Roma*
Massimo Collino, *Torino*
Teresa Esposito, *Naples*
Antonia Follenzi, *Novara*
Tommasina Guglielmelli, *Turin*
Lidia Larizza, *Milano*
Fortunato Lonardo, *Benevento*
Marco Lucarelli, *Rome*
Laura Mandelli, *Bologna*
Pier Giuseppe Pelicci, *Milan*
Daniela Perotti, *Milano*
Antonio M Persico, *Rome*
Enrico Pola, *Rome*
Giovanni Romeo, *Bologna*
Emanuela Signori, *Rome*
Alessio Squassina, *Cagliari*
Anna Stornaiuolo, *Milano*

**Japan**

Yutaka Hata, *Tokyo*
Mizukami Hiroaki, *Shimotsuke*
Johji Inazawa, *Tokyo*
Yasutomi Kamei, *Tokyo*
Tetsufumi Kanazawa, *Osaka*
Hiroyuki Kanzaki, *Miyagi*
Koichi Kawakami, *Shizuoka*
Akinori Kimura, *Tokyo*

Hiroshi Kunugi, *Tokyo*
Alexander Lezhava, *Kanagawa*
Hisao Masai, *Tokyo*
Ryuichi Morishita, *Suita*
Hitoshi Osaka, *Tochigi*
Masa-Aki Shibata, *Osaka*
Kazumi Suzukawa, *Tsukuba*
Masatoshi Tagawa, *Chiba*
Shin'ichi Takeda, *Tokyo*
Keizo Takenaga, *Shimane*
Hiroshi Tanooka, *Tokyo*
Takafumi Uchida, *Sendai*
Sho-ichi Yamagishi, *Kurume*
Toshiyuki Yamamoto, *Tokyo*
Norio Yasui-Furukori, *Hirosaki*

**Malaysia**

Ravindran Ankathil, *Kubang Kerian*
Hua Siew Gan, *Kubang Kerian*

**Mexico**

José F Munoz-Valle, *Zapopan*
Gilberto Vargas-Alarcon, *Mexico City*

**Netherlands**

Raoul CM Hennekam, *Amsterdam*
Anke H Maitland-van der Zee, *Utrecht*
Paulus J van Diest, *Utrecht*
Mannis van Oven, *Leiden*

**New Zealand**

Ryuji Fukuzawa, *Dunedin*
Christine M Morris, *Christchurch*

**Norway**

Kristian Tambs, *Oslo*
Martin Tesli, *Oslo*

**Pakistan**

Muhammad Naeem, *Islamabad*

**Peru**

Gustavo F Gonzales, *Lima*

**Poland**

Piotr Jan Wysocki, *Poznan*

**Portugal**

Manuel R Teixeira, *Porto*

**Russia**

Anton V Kiselev, *Saint-Petersburg*
Yuri B Yurov, *Moscow*

**Saudi Arabia**

Khaled K.S. Abu-Amero, *Riyadh*
Khawla S Al-Kuraya, *Riyadh*
Volodymyr Dvornyk, *Riyadh*
Wael Mohamed Swelam, *Medina*

**Serbia**

Lidija Radenovic, *Belgrade*

**Singapore**

N Gopalakrishna Iyer, *Singapore*
Peter E Lobie, *Singapore*
Eng-King Tan, *Singapore*

**Slovenia**

Borut Peterlin, *Ljubljana*
Uros Potocnik, *Maribor*

**South Korea**

Byung-Hoon Jeong, *Jeollabuk-do*
Jung Jin Kim, *Seoul*
Yonggoo Kim, *Seoul*
Taeg Kyu Kwon, *Taegu*
Seong-Wook Lee, *Yongin*
Taesung Park, *Seoul*
Han-Wook Yoo, *Seoul*

**Spain**

David Araújo-Vilar, *Santiago de Compostela*
Victor Asensi, *Oviedo*
Ignacio Blanco-Guillermo, *Barcelona*
Judit Sanz i Buxó, *Catalonia*
Javier S Castresana, *Pamplona*
Cristina Fillat, *Barcelona*
Karen Heath, *Madrid*
Adrián Llerena, *Badajoz*
Jose Felix Marti Masso, *San Sebastian*
José M Millán, *Valencia*
Salvador F Alino Pellilcer, *Valencia*
Juan Pie, *Zaragoza*
Jesus Prieto, *Pamplona*
Juan A Rey, *Madrid*
Miguel Chillon Rodriguez, *Bellaterra*
María E Sáez, *Seville*

**Sweden**

Mef Christina Nilbert, *Lund*

**Switzerland**

Angela Ciuffi, *Lausanne*
Daniel Schorderet, *Lausanne*

**Thailand**

Vorasuk Shotelersuk, *Bangkok*
Tewin Tencomnao, *Bangkok*
Viroj Wiwanitkit, *Bangkok*

**Turkey**

Mehmet Necdet Akkus, *Mersin*
Emrah Caylak, *Cankiri*
Merih Cetinkaya, *Bursa*
Ali Karaman, *Erzurum*
Mustafa Sahin, *Ankara*
Ahter Dilsad Sanlioglu, *Antalya*
Salih Sanlioglu, *Antalya*
Hülya Yazici, *Istanbul*

**Ukraine**

Ludmila Livshits, *Kyiv*

**United Arab Emirates**

Bassam R. Ali, *Al-Ain*

**United Kingdom**

Kristin Becker, *Cardiff*
Constantinos Demonacos, *Manchester*
David Jeremy Galton, *London*
Elaine Green, *Cardiff*
Wen G. Jiang, *Cardiff*
Patrick J Morrison, *Belfast*
Elisabeth Nacheva, *London*
Mary Callaghan Nicol, *Cardiff*
David J Timson, *Brighton*
Eugene M Tulchinsky, *Leicester*

**United States**

Praveen R Arany, *Cambridge*
Nedal Arar, *San Antonio*
Richard G Boles, *Los Angeles*
Merlin G Butler, *Kansas City*
Haifeng Chen, *Hayward*
J Don Chen, *Piscataway*
James J Chen, *Jefferson City*
Xiang-Ning Chen, *Richmond*
Qi Dai, *Nashville*
Shuo Dong, *Houston*
Yao-Shan Fan, *Miami*
Bing-Liang Fang, *Houston*
Peter J Francis, *Portland*
Xiaoyi Gao, *Chicago*
Yu-Bin Ge, *Detroit*
Antonio Giordano, *Philadelphia*

Thomas J Giordano, *Ann Arbor*
Stephen J Glatt, *Syracuse*
W Terrance Godbey, *New Orleans*
Dennis R Grayson, *Chicago*
Dong-Sheng Gu, *Indianapolis*
Zong-Sheng Guo, *Pittsburgh*
Hakon Hakonarson, *Philadelphia*
Wayne W Hancock, *Philadelphia*
David W Hein, *Louisville*
Huixiao Hong, *Jefferson*
Ji-Fan Hu, *Palo Atlo*
Shile Huang, *Shreveport*
Ying Huang, *Syracuse*
Johnny Huard, *Pittsburgh*
Barbara H Iglewski, *Rochester*
Yurij Ionov, *Buffalo*
Berit Kerner, *Los Angeles*
Alisa E Koch, *Ann Arbor*
Paul C Kuo, *Maywood*
Robert R Langley, *Houston*
Eduardo C Lau, *Milwaukee*
Kin-Hing W Lau, *Loma Linda*
Mong-Hong Lee, *Houston*
Da-Wei Li, *New Haven*
Shi-Bo Li, *Oklahoma City*
Wei Li, *Cleveland*
Ming-Fong Lin, *Omaha*
Steven R Lindheim, *Cincinnati*
Feng Liu, *Chapel Hill*
Xiao-Qi Liu, *West Lafayette*
Yao-Zhong Liu, *New Orleans*
Yong-Jun Liu, *New Orleans*
Bo Lu, *Nashville*
Qun Lu, *Greenville*
Xing-Guang Luo, *West Haven*
Richard C McEachin, *Ann Arbor*
Jian-Feng Meng, *Kansas*
Duane A Mitchell, *Durham*
Yin-Yuan Mo, *Springfield*
Viktor Morozov, *Bethesda*
Srinivas Mummidi, *San Antonio*
Mayumi Naramura, *Omaha*
Swapan K Nath, *Oklahoma*
Muthu Periasamy, *Columbus*
Gwendolyn P Quinn, *Tampa*
Rajalingam Raja, *Los Angeles*
Rajagopal Ramesh, *Houston*
Jasti S Rao, *Peoria*
Charles J Rosser, *Orlando*
Mark A Rothstein, *Louisville*
Ananda L Roy, *Boston*
Dharambir K Sanghera, *Oklahoma City*
Thomas Scholl, *Westborough*
Rong Shao, *Springfield*
Yuenian Eric Shi, *Roslyn Heights*
Shree Ram Singh, *Frederick*
Si-Hong Song, *Seminole*
Manjunath N Swamy, *El Paso*
Ming Tan, *Mobile*
Flora Tassone, *Sacramento*
Mustafa Tekin, *Miami*
Brad Therrell, *Austin*
Barry Trink, *Baltimore*
Tibor Valyi-Nagy, *Chicago*
Andre J van Wijnen, *Worcester*
Chia-Yeng Wang, *Chicago*
Jean Yin Jen Wang, *La Jolla*
Pin Wang, *Los Angeles*
Qing-Yi Wei, *Houston*
Qi-Ze Wei, *Manhattan*
David A Weinstein, *Gainesville*

Mary E Wilson, *Iowa City*
Xiaoling Xuei, *Indianapolis*
Guang Yang, *Augusta*
Zeng-Quan Yang, *Detroit*

Lin Yao, *Augusta*
Si-Ming Zhang, *Albuquerque*
Xiaoliu S Zhang, *Houston*
Tong Zhu, *Durham*



Uruguay

Jose L Badano, *Montevideo*



CASE REPORT

- 17 Mutation in *TNXB* gene causes moderate to severe Ehlers-Danlos syndrome
Kaufman CS, Butler MG

ABOUT COVER

Editorial Board Member of *World Journal of Medical Genetics*, Volodymyr Dvornyk, PhD, Professor, College of Science and General Studies, Alfaisal University, Riyadh 11533, Saudi Arabia

AIM AND SCOPE

World Journal of Medical Genetics (*World J Med Genet*, *WJMG*, online ISSN 2220-3184, DOI: 10.5496) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJMG covers topics concerning genes and the pathology of human disease, molecular analysis of simple and complex genetic traits, cancer genetics, epigenetics, gene therapy, developmental genetics, regulation of gene expression, strategies and technologies for extracting function from genomic data, pharmacological genomics, genome evolution. The current columns of *WJMG* include editorial, frontier, diagnostic advances, therapeutics advances, field of vision, mini-reviews, review, topic highlight, medical ethics, original articles, case report, clinical case conference (Clinicopathological conference), and autobiography.

We encourage authors to submit their manuscripts to *WJMG*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

World Journal of Medical Genetics is currently no indexing/abstracting.

FLYLEAF

I-IV Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Ya-Jing Lu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Fang-Fang Ji*
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL

World Journal of Medical Genetics

ISSN

ISSN 2220-3184 (online)

LAUNCH DATE

December 27, 2011

FREQUENCY

Quarterly

EDITOR-IN-CHIEF

Hans van Bokhoven, Professor, PhD, Department of Human Genetics and Cognitive Neurosciences, Radboud university Nijmegen Medical centre, PO Box 9101, 6500 HB Nijmegen, The Netherlands

EDITORIAL OFFICE

Jin-Lei Wang, Director
Xiu-Xia Song, Vice Director
World Journal of Medical Genetics

Room 903, Building D, Ocean International Center, No. 62 Dongsihuan Zhonglu, Chaoyang District, Beijing 100025, China
Telephone: +86-10-59080039
Fax: +86-10-85381893
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.wjgnet.com/esp/helpdesk.aspx>
<http://www.wjgnet.com>

PUBLISHER

Baishideng Publishing Group Inc
8226 Regency Drive,
Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.wjgnet.com/esp/helpdesk.aspx>
<http://www.wjgnet.com>

PUBLICATION DATE

May 27, 2016

COPYRIGHT

© 2016 Baishideng Publishing Group Inc. Articles published by this Open Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS

Full instructions are available online at http://www.wjgnet.com/bpg/g_info_20160116143427.htm

ONLINE SUBMISSION

<http://www.wjgnet.com/esp/>

Mutation in *TNXB* gene causes moderate to severe Ehlers-Danlos syndrome

Carolyn S Kaufman, Merlin G Butler

Carolyn S Kaufman, Merlin G Butler, Department of Psychiatry, Behavioral Sciences and Pediatrics, University of Kansas Medical Center, Kansas City, KS 66160, United States

Author contributions: Kaufman CS and Butler MG designed the report and wrote the manuscript.

Supported by The National Institute of Child Health and Human Development (NICHD), No. HD02528.

Institutional review board statement: University of Kansas Medical Center Institutional Review Board.

Informed consent statement: Verbal consent was obtained from the patient. There are no identifying images included in the manuscript.

Conflict-of-interest statement: There are no conflicts of interest.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Merlin G Butler, MD, PhD, Departments of Psychiatry, Behavioral Sciences and Pediatrics, University of Kansas Medical Center, 3901 Rainbow Blvd, MS 4015, Kansas City, KS 66160, United States. mbutler4@kumc.edu
Telephone: +1-913-5881800
Fax: +1-913-5881305

Received: April 16, 2016

Peer-review started: April 19, 2016

First decision: May 10, 2016

Revised: May 15, 2016

Accepted: May 18, 2016

Article in press: May 20, 2016

Published online: May 27, 2016

Abstract

We report a 28-year-old female who presented with severe joint pain, chronic muscle weakness, Raynaud's phenomenon, and hypermobility. She was found to have a 6074A > T nucleotide transition in the *TNXB* gene causing an amino acid protein change at Asp2025Val classified as likely pathogenic. We add this clinical report to the literature and classical human disease gene catalogs to identify this specific mutation as disease-causing. This gene variant was reported previously in a different 36-year-old patient who shared our patient's symptoms of joint hypermobility, skeletal and joint pain, skin elasticity and musculoskeletal problems, thereby causing a more severe presentation than seen in the hypermobility type of Ehlers-Danlos syndrome (EDS). At the time of writing, a few mutations in the *TNXB* gene have been recognized as pathogenic causing EDS due to tenascin-X deficiency, but the variant identified in our patient has not been recognized as pathogenic in online genetic databases. Our case study in combination with peer-reviewed literature suggests that the 6074A > T nucleotide transition in the *TNXB* gene may be classified as disease-causing for EDS due to tenascin-X deficiency.

Key words: Ehlers-Danlos syndrome; Genetic variants; Mutations; Hypermobility; Joint pain; Muscle weakness; Raynaud's phenomenon; Tenascin-X; *TNXB*; Ehlers-Danlos syndrome due to tenascin-X deficiency

© **The Author(s) 2016.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Various types of Ehlers-Danlos syndrome (EDS) have unique phenotypic features and genetic causes that are under investigation. This case report presents a gene variant (6074A > T nucleotide transition in the *TNXB* gene) not previously classified as disease-causing which we propose should be classified as pathogenic. This variant appears to produce joint hypermobility, skeletal pain, and musculoskeletal problems and should be classified as causing EDS due to tenascin-X deficiency.

Kaufman CS, Butler MG. Mutation in *TNXB* gene causes moderate to severe Ehlers-Danlos syndrome. *World J Med Genet* 2016; 6(2): 17-21 Available from: URL: <http://www.wjgnet.com/2220-3184/full/v6/i2/17.htm> DOI: <http://dx.doi.org/10.5496/wjmg.v6.i2.17>

INTRODUCTION

Ehlers-Danlos syndrome (EDS) encompasses a heterogeneous collection of connective tissue disorders with joint, skin and vascular involvement. Patients diagnosed with EDS have diverse clinical findings, which may be better understood and classified by identifying the genetic contribution to symptoms^[1,2]. Although this condition has been recognized since 1901^[3] and is now classified into nine types, delineation of the types and relationships to causative genes and their variants are underway, impacted by advances in genetic technology and next generation (exome) sequencing^[4,5]. Here, we assert that mutations in the *TNXB* gene, which encodes the glycoprotein tenascin-XB and related to the benign hypermobility form of EDS, may cause a more severe form or presentation of EDS.

CASE REPORT

Our 28-year-old patient showed no developmental delays as a child. She did undergo a right knee arthroscopic surgery not related to an injury at 18 years of age and breast augmentation at 21 years of age. She has had two pregnancies and was anemic during both pregnancies, due to decreased red blood cell count. The pregnancies were full-term and labor and delivery were normal. She experienced increased joint hypermobility during both of her pregnancies. Previous lab work showed normal thyroid and hepatic function. The patient had generalized joint pain since childhood with mild scoliosis. Anti-inflammatory agents did not relieve her pain. She has experienced instability of the right shoulder, pelvis and both hip joints with more problems with advancing age. During the clinic visit, we completed an assessment of the patient's joint mobility using the Beighton scale^[6], which results in a numerical score from 0 to 9 with one point assigned for the ability to perform each of the following actions: Passive dorsiflexion of the little finger beyond 90 degrees; passive apposition of the thumb to the flexor aspects of the forearm; hyperextension of the elbow beyond 10 degrees; hyperextension of the knee beyond 10 degrees; and forward flexion of the trunk, with knees straight, so that palms of hands rest easily on the floor. Our patient's Beighton score was 6 out of 9, including easy placement of palms to the floor (one point), hyperextension of both knees (two points), hyperextension of one elbow (one point), and passive apposition of both thumbs to the forearms (two points) as shown in Figure 1.

Furthermore, our patient also experienced Raynaud's

phenomenon, manifested as paresthesia with decreased blood flow in her fingers, lip, ears, and nose (Figure 1). This clinical presentation led to rheumatology evaluation including antibody studies and SED rate to diagnose a mixed connective tissue disorder such as lupus or rheumatoid arthritis. The results were normal. The patient's back pain continues to worsen and she experiences weakness throughout much of her body. She is unable to participate in mildly strenuous manual labor or movement related activities. Our patient identifies chronic pain and generalized muscle weakness as the most significant medical problems detrimental to daily functioning and quality of life.

The family history was limited due to little to no contact with her siblings and parents, all of whom live in Israel. Our patient has 100% Ashkenazi Jewish ancestry but has not had any Jewish specific genetic disease testing. She has a son who is 4 years of age and a daughter who is 10 years of age, both currently without relevant medical concerns or health problems.

We ordered next generation DNA sequencing panel and deletion/duplication analysis of 33 genes related to connective tissue disorders (*ABCC6*, *ACTA2*, *ATP7A*, *CBS*, *CHST14*, *COL1A1*, *COL1A2*, *COL2A1*, *COL3A1*, *COL5A1*, *COL5A2*, *COL9A1*, *COL9A2*, *COL9A3*, *COL11A1*, *COL11A2*, *ELN*, *FBN1*, *FBN2*, *FKBP14*, *FLNA*, *MYH11*, *MYLK*, *PLOD1*, *SKI*, *SLC2A10*, *SLC39A13*, *SMAD3*, *SMAD4*, *TGFB2*, *TGFBR1*, *TGFBR2*, and *TNXB*) performed commercially at the University of Nebraska Medical Center (Omaha, NE) using a blood sample from our patient. The only gene variant identified was a 6074A > T transition in the *TNXB* gene causing a Asp2025Val amino acid change at the protein level, classified as a variant of uncertain clinical significance. No known pathogenic deletions or duplications were detected within these genes.

DISCUSSION

Classification for EDS began in the late 1960s^[4] and continues to evolve and expand over time, as diagnoses move beyond focusing solely on clinical symptoms to a more genomics-driven approach. A popular simplified classification based on the Villenfranche nosology outlined the following six major types of EDS^[4]: Classical (Type I and Type II) with defect in type 5 collagen; Hypermobile (Type III) with unknown genetic causation; Vascular (Type IV) with abnormal type 3 collagen; Kyphoscoliosis (Type VI) with deficiency of lysyl hydroxylase (LH); Arthrochalasia (Type VII) with deficiency of chains in type I collagen; and Dermatosparaxis (Type VIII) with deficiency of procollagen N-proteinase enzyme in type I collagen.

In a more recent review^[5], this classification was expanded based on current evidence, and a total of nine types of EDS were outlined: Classical (caused by defects in collagen *COL5A1* and *COL5A2* genes) involving skin laxity, joint hypermobility, widened scars, and easy bruising; Hypermobile (unknown cause) with velvet




Figure 1 Patient displays extreme hyperflexibility as represented by bilateral positive thumb to wrist signs on the Beighton hyperflexibility scale^[6]. Raynaud's phenomenon of the fingers with cool temperature and decreased blood flow.

skin and laxity, joint dislocations and hypermobility; Vascular (*COL3A1* gene) with arterial rupture, hypermobility, varicose veins, thin translucent skin, and easy bruising; Kyphoscoliosis (LH 1 - *PLOD1* gene) with muscle hypotonia in infants, scoliosis from birth, joint laxity, gross motor delay, easy bruising, and fragility of sclerae; Arthrochalasia (collagen 1 - *COL1A1*, *COL1A2* genes) with severe joint hypermobility, congenital bilateral hip dislocations, skin elasticity, easy bruising, kyphoscoliosis, and osteopenia; Dermatosparaxis (pro-collagen N-proteinase - *ADAMTS2* gene) with sagging redundant skin, blue sclerae, short stature, and umbilical hernia; EDS with scoliosis, myopathy, hearing impairment (*FKBP14* gene); Musculocontractural EDS (*CHST14* gene) with progressive kyphoscoliosis, adducted thumbs in infancy, joint hypermobility, clubfoot, arachnodactyly, elastic skin, and poor wound healing; and finally Tenascin-X deficient (*TNXB* gene). Those with Tenascin-X (*TNXB*) deficient EDS involve features of marked skin laxity, pronounced joint hypermobility and severe bruising^[5]. We extend the description of those with mutations in the *TNXB* gene to include significant joint pain and Raynaud's phenomenon, thus resulting in a more severe form of EDS than previously classified. Whether the severity depends on the type or location of the gene variant (mutation) is unknown.

The *TNXB* gene (sometimes known by other names such as *HXBL*, *TNX*, *TENX*, *XB*) belongs to a family of genes called fibronectin type III domain containing and localizes to the major histocompatibility complex

(MHC) class III region on chromosome 6, at cytogenetic location 6p21.3. The *TNXB* gene encodes for a protein called tenascin-X, which may regulate assembly and production of specific types of collagen. Tenascin-X helps organize and maintain the structure of tissues supporting the joints, organs, skin, connective tissues, and muscles^[7].

Tenascin-X is an extracellular matrix glycoprotein, thought to function in matrix maturation during wound healing^[7] and became the first *EDS* gene that does not encode a collagen-modifying enzyme or a fibrillar collagen^[8]. Mice knockouts (*Tnxb*^{-/-}) showed skin hyperextensibility, reflecting the human phenotype of EDS, and the skin collagen content was significantly reduced compared to wildtype. Additionally, the density of fibrils in the skin of knockout (*Tnxb*^{-/-}) mice was reduced compared to wildtype, suggesting tenascin-X plays a significant regulatory role in collagen deposition by dermal fibroblasts^[8]. Hence, tenascin-X appears to be involved in the regulation of the structure and stability of elastic fibers, which give elasticity and flexibility to connective tissues.

Alterations in the *TNXB* gene and effects on protein levels have also been studied^[9]. The researchers measured tenascin-X in serum from 21 healthy subjects and 151 patients diagnosed with EDS. They reported that 146 of the 151 patients with EDS had detectable tenascin-X levels in their serum. A closer examination of the 5 patients diagnosed with EDS and lacking serum tenascin-X revealed that all had hyperelastic skin and

hypermobility, and a subset had joint pain and multiple subluxations. One patient was homozygous and another heterozygous for a 30-kb deletion in the *TNXB* gene resulting in a nonfunctional fusion gene. Two patients were homozygous for a 2-bp deletion in exon 8 of the *TNXB* gene, which altered the open reading frame coding for amino acids 1184 through 1230 before causing a premature stop codon. The fifth patient was homozygous for a 2-bp insertion in exon 3 which caused a stop codon at position 707. These genetic variations are now classified as causing EDS specifically due to tenascin-X protein deficiency.

Furthermore, a different study^[10] identified three missense mutations in the *TNXB* gene, one in each of three patients diagnosed with the hypermobility type EDS but with normal tenascin-X serum levels: A 3582A > G transition (Val1195Met), an 85C > T transition (Arg29Trp), and a 12097C > A transition (Leu4033Ile). These wild type amino acids and locations are strongly conserved in this protein. The researchers then compared skin biopsies from these patients with other EDS hypermobility type patients who did not have gene variants in the *TNXB* gene and determined that elastic fiber abnormalities in hypermobility type EDS are specific for TNX-haploinsufficient individuals and confirm an important role for TNX in regulating elastic fiber integrity in humans^[10].

As demonstrated in the literature by examples described, a variety of mutations, deletions, and insertions in the *TNXB* gene may cause features of EDS. Our patient's gene change was classified as a variant of unknown significance and not as yet classified as disease-causing. This missense mutation in exon 17 of the *TNXB* gene in our patient is moderately conserved with an in silico prediction of trending damaging. This mutation causes a significant amino acid change from an aspartic acid which is negatively charged to a valine which is non-polar; this transition could theoretically impact protein structure due to such a dramatic change in the amino acid classification type. Based on our patient's clinical symptoms and DNA findings with evidence from the literature, we propose that the 6074A > T nucleotide transition (p.Asp2025Val) identified in our patient causes a moderate to severe form of EDS that includes significant joint pain, muscle weakness, hypermobility and Raynaud's phenomenon.

A limitation of our report is the lack of contact by the patient with her siblings or parents, all of whom live in Israel. Therefore, it is not possible for us to reach members of her family to identify the presence or absence of the gene variant and to better characterize the genotype-phenotype relationship. Furthermore, our patient lives at a considerable distance from the medical center making travel difficult for her and her immediate family (e.g., younger children). It is unlikely that arrangements could be made in a timely fashion to undertake skin biopsy and assessment of elastic fibers in the patient or her living children without evidence of disease in which an invasive procedure is not warranted

even if accessible.

Our patient is also of Ashkenazi Jewish ancestry. A 36-year-old patient with this same amino acid change at the protein level reported to be likely pathogenic^[2,11-14] and that patient was diagnosed clinically with the hypermobility type of EDS. The patient's symptoms included hypermobility, skeletal pain, skin elasticity, gastrointestinal and musculoskeletal problems, all consistent with EDS. The status of Jewish ancestry with this reported patient is unknown. We propose that both of these patients may have a form of EDS due to their specific mutation in the *TNXB* gene, causing more severe symptoms sufficiently significant to interfere with daily functioning and quality of life. More research is needed to conclude that this gene variant is causative and authors encourage the reporting of other patients with variants of the *TNXB* gene, clinical features and genotype-phenotype correlations.

ACKNOWLEDGMENTS

We thank our patient.

COMMENTS

Case characteristics

A 28-year-old female presented with joint pain, muscle weakness, hypermobility, and Raynaud's phenomenon. She was referred to our clinic for evaluation for a connective tissue disorder.

Clinical diagnosis

Severe joint pain, chronic muscle weakness, Raynaud's phenomenon, hypermobility with Beighton hypermobility score of 6 out of 9, mild scoliosis.

Differential diagnosis

Ehlers-Danlos syndrome.

Laboratory diagnosis

Previous rheumatology evaluation ruled out lupus or rheumatoid arthritis through antibody studies and SED rate. Testing through the University of Nebraska Medical Center (Omaha, NE) showed a nucleotide 6074A > T transition in the *TNXB* gene using a blood sample.

Treatment

Reduce daily stress on joints by minimizing mildly strenuous activity. Avoid hyper-extending joints.

Related report

Lee H, Deignan JL, Dorrani N, Strom SP, Kantarci S, Quintero-Rivera F, Das K, Toy T, Harry B, Yourshaw M, Fox M, Fogel BL, Martinez-Agosto JA, Wong DA, Chang VY, Shieh PB, Palmer CG, Dipple KM, Grody WW, Vilain E, Nelson SF. Clinical exome sequencing for genetic identification of rare Mendelian disorders. *JAMA* 2014; 312: 1880-1887 [PMID: 25326637 DOI: 10.1001/jama.2014.14604].

Term explanation

Ehlers-Danlos syndrome includes a heterogeneous collection of connective tissue disorders with vascular, skeletal and skin involvement.

Experiences and lesson

As technology advances, authors are able to order more sophisticated genetic testing in the clinical setting, such as the sequencing and deletion/duplication panel ordered in this case report. However, the meaning of the results is not always clear, as more often than not they identify "variants of unknown clinical

significance", meaning the change at the DNA level does not have a clear, recognized outcome for the phenotype. In these cases, they use computer simulation to predict whether or not the change may harmful to protein structure or function. They also review literature for cases involving the variant, with the hope of associating genotype with phenotype. In the case of the patient reported in this manuscript, they concluded through in-depth examination of the variant and a comprehensive literature review that the 6074A > T transition in the *TNXB* gene seen in our patient is likely pathogenic and causes a moderate to severe form of Ehlers-Danlos syndrome.

Peer-review

The authors report a case of Ehlers-Danlos syndrome with a mutation in *TNXB* gene. The report is interesting.

REFERENCES

- 1 **Online Mendelian Inheritance in Man.** Johns Hopkins University, Baltimore, MD. [updated 2014 Aug 26]. Available from: URL: <http://omim.org/entry/606408?search=tnxb&highlight=tnxb>
- 2 **Lee H,** Deignan JL, Dorrani N, Strom SP, Kantarci S, Quintero-Rivera F, Das K, Toy T, Harry B, Yourshaw M, Fox M, Fogel BL, Martinez-Agosto JA, Wong DA, Chang VY, Shieh PB, Palmer CG, Dipple KM, Grody WW, Vilain E, Nelson SF. Clinical exome sequencing for genetic identification of rare Mendelian disorders. *JAMA* 2014; **312**: 1880-1887 [PMID: 25326637 DOI: 10.1001/jama.2014.14604]
- 3 **Ehlers E.** Cutis laxa, neigung zu harmorrhagien in der haut, lockerun mehrer Artikulationen. *Dermat Ztschr* 1901; **8**: 173
- 4 **Parapia LA,** Jackson C. Ehlers-Danlos syndrome--a historical review. *Br J Haematol* 2008; **141**: 32-35 [PMID: 18324963 DOI: 10.1111/j.1365-2141.2008.06994.x]
- 5 **Sobey G.** Ehlers-Danlos syndrome: how to diagnose and when to perform genetic tests. *Arch Dis Child* 2015; **100**: 57-61 [PMID: 24994860 DOI: 10.1136/archdischild-2013-304822]
- 6 **Beighton P,** Solomon L, Soskolne CL. Articular mobility in an African population. *Ann Rheum Dis* 1973; **32**: 413-418 [PMID: 4751776 DOI: 10.1136/ard.32.5.413]
- 7 **National Library of Medicine (US).** Genetics Home Reference [Internet]. Bethesda (MD): The Library; 2016 Mar 28. TNXB. [reviewed 2015 Nov; cited 2016 Apr 7]. Available from: URL: <http://ghr.nlm.nih.gov/gene/TNXB>
- 8 **Mao JR,** Taylor G, Dean WB, Wagner DR, Afzal V, Lotz JC, Rubin EM, Bristow J. Tenascin-X deficiency mimics Ehlers-Danlos syndrome in mice through alteration of collagen deposition. *Nat Genet* 2002; **30**: 421-425 [PMID: 11925569 DOI: 10.1038/ng850]
- 9 **Schalkwijk J,** Zweers MC, Steijlen PM, Dean WB, Taylor G, van Vlijmen IM, van Haren B, Miller WL, Bristow J. A recessive form of the Ehlers-Danlos syndrome caused by tenascin-X deficiency. *N Engl J Med* 2001; **345**: 1167-1175 [PMID: 11642233 DOI: 10.1056/NEJMoa002939]
- 10 **Zweers MC,** Dean WB, van Kuppevelt TH, Bristow J, Schalkwijk J. Elastic fiber abnormalities in hypermobility type Ehlers-Danlos syndrome patients with tenascin-X mutations. *Clin Genet* 2005; **67**: 330-334 [PMID: 15733269 DOI: 10.1111/j.1399-0004.2005.00401.x]
- 11 **Beighton P,** Horan F. Orthopaedic aspects of the Ehlers-Danlos syndrome. *J Bone Joint Surg Br* 1969; **51**: 444-453 [PMID: 5820785]
- 12 **Beighton P,** De Paepe A, Steinmann B, Tsipouras P, Wenstrup RJ. Ehlers-Danlos syndromes: revised nosology, Villefranche, 1997. Ehlers-Danlos National Foundation (USA) and Ehlers-Danlos Support Group (UK). *Am J Med Genet* 1998; **77**: 31-37 [PMID: 9557891 DOI: 10.1002/(SICI)1096-8628(19980428)77:1<31::AID-AJMG8>3.0.CO;2-O]
- 13 **De Paepe A,** Malfait F. The Ehlers-Danlos syndrome, a disorder with many faces. *Clin Genet* 2012; **82**: 1-11 [PMID: 22353005 DOI: 10.1111/j.1399-0004.2012.01858.x]
- 14 **Juul-Kristensen B,** Røgind H, Jensen DV, Remvig L. Inter-examiner reproducibility of tests and criteria for generalized joint hypermobility and benign joint hypermobility syndrome. *Rheumatology (Oxford)* 2007; **46**: 1835-1841 [PMID: 18006569 DOI: 10.1093/rheumatology/kem290]

P- Reviewer: Castori M, Sakamoto A, Sakkas LI **S- Editor:** Qiu S
L- Editor: A **E- Editor:** Lu YJ





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>



World Journal of *Medical Genetics*

World J Med Genet 2016 August 27; 6(3): 22-33





Editorial Board

2016-2019

The *World Journal of Medical Genetics* Editorial Board consists of 331 members, representing a team of worldwide experts in medical genetics. They are from 50 countries, including Argentina (1), Armenia (1), Australia (4), Austria (2), Belgium (2), Brazil (6), Bulgaria (2), Canada (6), Chile (1), China (39), Croatia (1), Czech Republic (1), Denmark (3), Ecuador (1), Egypt (2), France (10), Germany (4), Greece (2), Hungary (1), India (13), Iran (7), Ireland (1), Israel (2), Italy (22), Japan (23), Malaysia (2), Mexico (2), Netherlands (5), New Zealand (2), Norway (2), Pakistan (1), Peru (1), Poland (1), Portugal (1), Russia (2), Saudi Arabia (4), Serbia (1), Singapore (3), Slovenia (2), South Korea (7), Spain (16), Sweden (1), Switzerland (2), Thailand (3), Turkey (8), Ukraine (1), United Arab Emirates (1), United Kingdom (10), United States (95) and Uruguay (1).

EDITOR-IN-CHIEF

Hans van Bokhoven, *Nijmegen*

GUEST EDITORIAL BOARD MEMBERS

Chia-Hsiang Chen, *Zhunan*
Ji-Yih Chen, *Keelung*
Bor-Luen Chiang, *Taipei*
Yu-Chen Hu, *Hsinchu*
Suh-Hang H Juo, *Kaohsiung*
Hsien-Hsiung Lee, *Taichung*
Yueh-Lun Lee, *Taipei*
Chin-San Liu, *Changhua*
Pei-Jung Lu, *Tainan*
Chien-Kuo Tai, *Chiayi*
Po-Nien Tsao, *Taipei*
Wendy Wen-Hui Weng, *Taipei*
Kuo-Liang Yang, *Hualien*

MEMBERS OF THE EDITORIAL BOARD



Argentina

Carlos J Pirola, *Buenos Aires*



Armenia

Hasmik Mkrtchyan, *Yerevan*



Australia

Simon Easteal, *Canberra*
Jeremy BM Jowett, *Melbourne*

Karen Anne Mather, *Randwick*
Grant Morahan, *Perth*



Austria

Harald N Aschauer, *Vienna*
Anton Wutz, *Vienna*



Belgium

Teresinha Leal, *Brussels*
van West, Dirk, *Antwerp*



Brazil

Pereira Priscila Krauss, *Rio de Janeiro*
Sharbel Weidner Maluf, *Porto Alegre*
Milton Ozório Moraes, *Rio de Janeiro*
Fernando Regla Vargas, *Rio de Janeiro*
Jeane Eliete Laguila Visentainer, *Sao Paulo*
Isabela Jube Wastowski, *Goiania*



Bulgaria

Spaska Angelova Stanilova, *Stara Zagora*
Albena Todorova, *Sofia*



Canada

Régen Drouin, *Sherbrooke*
Gonzalo Hortelano, *Hamilton*
William Jia, *Vancouver*

Sean Li, *Ottawa*
Shiva M Singh, *London*
Liting Song, *Toronto*



Chile

Jara Sosa Lilian Elena, *Santiago*



China

George G Chen, *Hong Kong*
Ning-Han Feng, *Nanjing*
Shi-Ang Huang, *Wuhan*
Wei Huang, *Shanghai*
Chun-Yan Ji, *Jinan*
Hong-Chuan Jin, *Hangzhou*
Ke-Shen Li, *Zhanjiang*
Zhuo-Zhuang Lu, *Beijing*
Xiao-Ping Luo, *Wuhan*
Wen-Bin Qian, *Hangzhou*
Rong Rui, *Nanjing*
Lun-Quan Sun, *Changsha*
Qian Tao, *Hong Kong*
George Sai-Wah Tsao, *Hong Kong*
Ming-Rong Wang, *Beijing*
Shi-Jie Wang, *Shijiazhuang*
Wei Wang, *Beijing*
Ji Wu, *Shanghai*
Hua-Xi Xu, *Zhenjiang*
Ruian Xu, *Xiamen*
Wei-Hua Yan, *Linhai*
Rong-Cun Yang, *Tianjin*
Bang-Ce Ye, *Shanghai*
Rui-Xing Yin, *Nanning*
Yi Zeng, *Beijing*
Xiao-Feng Zhu, *Guangzhou*

**Croatia**

Gordan Lauc, *Zagreb*

**Czech Republic**

Marie Lipoldová, *Prague*

**Denmark**

Vibeke Andersen, *Odense*
Thomas V. O. Hansen, *Copenhagen*
Goutam Sahana, *Tjele*

**Ecuador**

Cesar Paz-Y-Mino, *Quito*

**Egypt**

Menha MA Swellam, *Giza*
Samia Ali Temtamy, *Cairo*

**France**

Christophe Chevillard, *Marseille*
Johanna Chluba, *Dijon*
Zdenko Herceg, *Lyon*
Enzo Lalli, *Valbonne*
James Lespinasse, *Chambéry*
Bernard S. Lopez, *Fontenay aux Roses*
Eric Pasmant, *Clichy*
Georges Vassaux, *Nice Cedex*
Kay-Dietrich Wagner, *Nice Cedex*
Qing Wang, *Lyon*

**Germany**

Stefan Bohringer, *Essen*
Anibh Martin Das, *Hannover*
Gudrun A. Rappold, *Heidelberg*
Hermona Soreq, *Jerusalem*

**Greece**

Roxani Angelopoulou, *Athens*
Voula Velissariou, *Athens*

**Hungary**

Béla Meleg, *Pecs*

**India**

Arvind Kumar Arya, *Meerut*
Prakash Sadashiv Gambhir, *Pune*
Katta KM Girisha, *Karnataka*
KAISER Jamil, *Hyderabad*

Narendra Joshi, *Navi Mumbai*
Swarna Mandava, *Mumbai*
Chinmay K Panda, *Kolkata*
Rajni Rani, *New Delhi*
Susanta Roychoudhury, *Kolkata*
Sudeep R Shah, *Mumbai*
Ranbir Chander Sobti, *Chandigarh*
Meera Vaswani, *New Delhi*
Suresh P Vyas, *Sagar*

**Iran**

Yahya Daneshbod, *Shiraz*
Fariborz Ghaffarpasand, *Shiraz*
Mohammad H Modarressi, *Tehran*
Shirin Hasani Ranjbar, *Tehran*
Nader tajik, *Tehran*
Dor Mohammad Kordi Tamandani, *Zahedan*
Alireza Zomorodipour, *Tehra*

**Ireland**

Ross McManus, *Dublin*

**Israel**

Aliza Shlomit Amiel, *Kfar-Saba*
Rachel Ben-Shlomo, *Tivon*

**Italy**

Francesco Acquati, *Varese*
Nenad Bukvic, *Bari*
Gabriele Candiani, *Milan*
Antonio Cao, *Cagliari*
Raffaella Cascella, *Rome*
Marco Castori, *Roma*
Massimo Collino, *Torino*
Teresa Esposito, *Naples*
Antonia Follenzi, *Novara*
Tommasina Guglielmelli, *Turin*
Lidia Larizza, *Milano*
Fortunato Lonardo, *Benevento*
Marco Lucarelli, *Rome*
Laura Mandelli, *Bologna*
Pier Giuseppe Pelicci, *Milan*
Daniela Perotti, *Milano*
Antonio M Persico, *Rome*
Enrico Pola, *Rome*
Giovanni Romeo, *Bologna*
Emanuela Signori, *Rome*
Alessio Squassina, *Cagliari*
Anna Stornaiuolo, *Milano*

**Japan**

Yutaka Hata, *Tokyo*
Mizukami Hiroaki, *Shimotsuke*
Johji Inazawa, *Tokyo*
Yasutomi Kamei, *Tokyo*
Tetsufumi Kanazawa, *Osaka*
Hiroyuki Kanzaki, *Miyagi*
Koichi Kawakami, *Shizuoka*
Akinori Kimura, *Tokyo*

Hiroshi Kunugi, *Tokyo*
Alexander Lezhava, *Kanagawa*
Hisao Masai, *Tokyo*
Ryuichi Morishita, *Suita*
Hitoshi Osaka, *Tochigi*
Masa-Aki Shibata, *Osaka*
Kazumi Suzukawa, *Tsukuba*
Masatoshi Tagawa, *Chiba*
Shin'ichi Takeda, *Tokyo*
Keizo Takenaga, *Shimane*
Hiroshi Tanooka, *Tokyo*
Takafumi Uchida, *Sendai*
Sho-ichi Yamagishi, *Kurume*
Toshiyuki Yamamoto, *Tokyo*
Norio Yasui-Furukori, *Hirosaki*

**Malaysia**

Ravindran Ankathil, *Kubang Kerian*
Hua Siew Gan, *Kubang Kerian*

**Mexico**

José F Munoz-Valle, *Zapopan*
Gilberto Vargas-Alarcon, *Mexico City*

**Netherlands**

Raoul CM Hennekam, *Amsterdam*
Anke H Maitland-van der Zee, *Utrecht*
Paulus J van Diest, *Utrecht*
Mannis van Oven, *Leiden*

**New Zealand**

Ryuji Fukuzawa, *Dunedin*
Christine M Morris, *Christchurch*

**Norway**

Kristian Tambs, *Oslo*
Martin Tesli, *Oslo*

**Pakistan**

Muhammad Naeem, *Islamabad*

**Peru**

Gustavo F Gonzales, *Lima*

**Poland**

Piotr Jan Wysocki, *Poznan*

**Portugal**

Manuel R Teixeira, *Porto*



Russia

Anton V Kiselev, *Saint-Petersburg*
Yuri B Yurov, *Moscow*



Saudi Arabia

Khaled K.S. Abu-Amero, *Riyadh*
Khawla S Al-Kuraya, *Riyadh*
Volodymyr Dvornyk, *Riyadh*
Wael Mohamed Swelam, *Medina*



Serbia

Lidija Radenovic, *Belgrade*



Singapore

N Gopalakrishna Iyer, *Singapore*
Peter E Lobie, *Singapore*
Eng-King Tan, *Singapore*



Slovenia

Borut Peterlin, *Ljubljana*
Uros Potocnik, *Maribor*



South Korea

Byung-Hoon Jeong, *Jeollabuk-do*
Jung Jin Kim, *Seoul*
Yonggoo Kim, *Seoul*
Taeg Kyu Kwon, *Taegu*
Seong-Wook Lee, *Yongin*
Taesung Park, *Seoul*
Han-Wook Yoo, *Seoul*



Spain

David Araújo-Vilar, *Santiago de Compostela*
Victor Asensi, *Oviedo*
Ignacio Blanco-Guillermo, *Barcelona*
Judit Sanz i Buxó, *Catalonia*
Javier S Castresana, *Pamplona*
Cristina Fillat, *Barcelona*
Karen Heath, *Madrid*
Adrián Llerena, *Badajoz*
Jose Felix Marti Masso, *San Sebastian*
José M Millán, *Valencia*
Salvador F Alino Pellilcer, *Valencia*
Juan Pie, *Zaragoza*
Jesus Prieto, *Pamplona*
Juan A Rey, *Madrid*
Miguel Chillon Rodriguez, *Bellaterra*
María E Sáez, *Seville*



Sweden

Mef Christina Nilbert, *Lund*



Switzerland

Angela Ciuffi, *Lausanne*
Daniel Schorderet, *Lausanne*



Thailand

Vorasuk Shotelersuk, *Bangkok*
Tewin Tencomnao, *Bangkok*
Viroj Wiwanitkit, *Bangkok*



Turkey

Mehmet Necdet Akkus, *Mersin*
Emrah Caylak, *Cankiri*
Merih Cetinkaya, *Bursa*
Ali Karaman, *Erzurum*
Mustafa Sahin, *Ankara*
Ahter Dilsad Sanlioglu, *Antalya*
Salih Sanlioglu, *Antalya*
Hülya Yazici, *Istanbul*



Ukraine

Ludmila Livshits, *Kyiv*



United Arab Emirates

Bassam R. Ali, *Al-Ain*



United Kingdom

Kristin Becker, *Cardiff*
Constantinos Demonacos, *Manchester*
David Jeremy Galton, *London*
Elaine Green, *Cardiff*
Wen G. Jiang, *Cardiff*
Patrick J Morrison, *Belfast*
Elisabeth Nacheva, *London*
Mary Callaghan Nicol, *Cardiff*
David J Timson, *Brighton*
Eugene M Tulchinsky, *Leicester*



United States

Praveen R Arany, *Cambridge*
Nedal Arar, *San Antonio*
Richard G Boles, *Los Angeles*
Merlin G Butler, *Kansas City*
Haifeng Chen, *Hayward*
J Don Chen, *Piscataway*
James J Chen, *Jefferson City*
Xiang-Ning Chen, *Richmond*
Qi Dai, *Nashville*
Shuo Dong, *Houston*
Yao-Shan Fan, *Miami*
Bing-Liang Fang, *Houston*
Peter J Francis, *Portland*
Xiaoyi Gao, *Chicago*
Yu-Bin Ge, *Detroit*
Antonio Giordano, *Philadelphia*

Thomas J Giordano, *Ann Arbor*
Stephen J Glatt, *Syracuse*
W Terrance Godbey, *New Orleans*
Dennis R Grayson, *Chicago*
Dong-Sheng Gu, *Indianapolis*
Zong-Sheng Guo, *Pittsburgh*
Hakon Hakonarson, *Philadelphia*
Wayne W Hancock, *Philadelphia*
David W Hein, *Louisville*
Huixiao Hong, *Jefferson*
Ji-Fan Hu, *Palo Atlo*
Shile Huang, *Shreveport*
Ying Huang, *Syracuse*
Johnny Huard, *Pittsburgh*
Barbara H Iglewski, *Rochester*
Yurij Ionov, *Buffalo*
Berit Kerner, *Los Angeles*
Alisa E Koch, *Ann Arbor*
Paul C Kuo, *Maywood*
Robert R Langley, *Houston*
Eduardo C Lau, *Milwaukee*
Kin-Hing W Lau, *Loma Linda*
Mong-Hong Lee, *Houston*
Da-Wei Li, *New Haven*
Shi-Bo Li, *Oklahoma City*
Wei Li, *Cleveland*
Ming-Fong Lin, *Omaha*
Steven R Lindheim, *Cincinnati*
Feng Liu, *Chapel Hill*
Xiao-Qi Liu, *West Lafayette*
Yao-Zhong Liu, *New Orleans*
Yong-Jun Liu, *New Orleans*
Bo Lu, *Nashville*
Qun Lu, *Greenville*
Xing-Guang Luo, *West Haven*
Richard C McEachin, *Ann Arbor*
Jian-Feng Meng, *Kansas*
Duane A Mitchell, *Durham*
Yin-Yuan Mo, *Springfield*
Viktor Morozov, *Bethesda*
Srinivas Mummidi, *San Antonio*
Mayumi Naramura, *Omaha*
Swapan K Nath, *Oklahoma*
Muthu Periasamy, *Columbus*
Gwendolyn P Quinn, *Tampa*
Rajalingam Raja, *Los Angeles*
Rajagopal Ramesh, *Houston*
Jasti S Rao, *Peoria*
Charles J Rosser, *Orlando*
Mark A Rothstein, *Louisville*
Ananda L Roy, *Boston*
Dharambir K Sanghera, *Oklahoma City*
Thomas Scholl, *Westborough*
Rong Shao, *Springfield*
Yuenian Eric Shi, *Roslyn Heights*
Shree Ram Singh, *Frederick*
Si-Hong Song, *Seminole*
Manjunath N Swamy, *El Paso*
Ming Tan, *Mobile*
Flora Tassone, *Sacramento*
Mustafa Tekin, *Miami*
Brad Therrell, *Austin*
Barry Trink, *Baltimore*
Tibor Valyi-Nagy, *Chicago*
Andre J van Wijnen, *Worcester*
Chia-Yeng Wang, *Chicago*
Jean Yin Jen Wang, *La Jolla*
Pin Wang, *Los Angeles*
Qing-Yi Wei, *Houston*
Qi-Ze Wei, *Manhattan*
David A Weinstein, *Gainesville*

Mary E Wilson, *Iowa City*
Xiaoling Xuei, *Indianapolis*
Guang Yang, *Augusta*
Zeng-Quan Yang, *Detroit*

Lin Yao, *Augusta*
Si-Ming Zhang, *Albuquerque*
Xiaoliu S Zhang, *Houston*
Tong Zhu, *Durham*



Uruguay

Jose L Badano, *Montevideo*



REVIEW

- 22 Chitosan DNA nanoparticles for oral gene delivery

Patel BJ, Vignesh NK, Hortelano G

Contents

World Journal of Medical Genetics
Volume 6 Number 3 August 27, 2016

ABOUT COVER

Editorial Board Member of *World Journal of Medical Genetics*, Sharbel Weidner Maluf, PhD, Doctor, Medical Genetics Service, Hospital de Clínicas de Porto Alegre, Porto Alegre 90035-903, Brazil

AIM AND SCOPE

World Journal of Medical Genetics (*World J Med Genet*, *WJMG*, online ISSN 2220-3184, DOI: 10.5496) is a peer-reviewed open access academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJMG covers topics concerning genes and the pathology of human disease, molecular analysis of simple and complex genetic traits, cancer genetics, epigenetics, gene therapy, developmental genetics, regulation of gene expression, strategies and technologies for extracting function from genomic data, pharmacological genomics, genome evolution. The current columns of *WJMG* include editorial, frontier, diagnostic advances, therapeutics advances, field of vision, mini-reviews, review, topic highlight, medical ethics, original articles, case report, clinical case conference (Clinicopathological conference), and autobiography.

We encourage authors to submit their manuscripts to *WJMG*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great basic and clinical significance.

INDEXING/ABSTRACTING

World Journal of Medical Genetics is currently no indexing/abstracting.

FLYLEAF

I-IV Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Ya-Jing Lu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Xue-Mei Gong*
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL

World Journal of Medical Genetics

ISSN

ISSN 2220-3184 (online)

LAUNCH DATE

December 27, 2011

FREQUENCY

Quarterly

EDITOR-IN-CHIEF

Hans van Bokhoven, Professor, PhD, Department of Human Genetics and Cognitive Neurosciences, Radboud university Nijmegen Medical centre, PO Box 9101, 6500 HB Nijmegen, The Netherlands

EDITORIAL OFFICE

Jin-Lei Wang, Director
Xiu-Xia Song, Vice Director
World Journal of Medical Genetics

Room 903, Building D, Ocean International Center,
No. 62 Dongsihuan Zhonglu, Chaoyang District,
Beijing 100025, China
Telephone: +86-10-59080039
Fax: +86-10-85381893
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>
<http://www.wjgnet.com>

PUBLISHER

Baishideng Publishing Group Inc
8226 Regency Drive,
Pleasanton, CA 94588, USA
Telephone: +1-925-223-8242
Fax: +1-925-223-8243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>
<http://www.wjgnet.com>

PUBLICATION DATE

August 27, 2016

COPYRIGHT

© 2016 Baishideng Publishing Group Inc. Articles published by this Open Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS

<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION

<http://www.wjgnet.com/esps/>

Chitosan DNA nanoparticles for oral gene delivery

Bhavika J Patel, Nithin K Vignesh, Gonzalo Hortelano

Bhavika J Patel, Nithin K Vignesh, Department of Health Sciences, McMaster University, Hamilton, Ontario L8N 3Z5, Canada

Gonzalo Hortelano, Department of Biology, School of Science and Technology, Nazarbayev University, Astana 010000, Kazakhstan

Gonzalo Hortelano, School of Biomedical Engineering, McMaster University, Hamilton, Ontario L8N 3Z5, Canada

Gonzalo Hortelano, Department of Pathology and Molecular Medicine, McMaster University, Hamilton, Ontario L8N 3Z5, Canada

Author contributions: All authors contributed to this paper with conception and design of the study, literature review and analysis, drafting and critical revision and editing, and final approval of the final version.

Conflict-of-interest statement: Non declared.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Correspondence to: Gonzalo Hortelano, PhD, Department of Biology, School of Science and Technology, Nazarbayev University, Astana 010000, Kazakhstan. gonzalo.hortelano@nu.edu.kz
Telephone: +7-717-2705782

Received: January 17, 2016

Peer-review started: January 18, 2016

First decision: February 29, 2016

Revised: July 20, 2016

Accepted: August 6, 2016

Article in press: August 8, 2016

Published online: August 27, 2016

Abstract

Gene therapy is a promising technology with potential applications in the treatment of medical conditions, both congenital and acquired. Despite its label as breakthrough technology for the 21st century, the simple concept of gene therapy - the introduction of a functional copy of desired genes in affected individuals - is proving to be more challenging than expected. Oral gene delivery has shown intriguing results and warrants further exploration. In particular, oral administration of chitosan DNA nanoparticles, one the most commonly used formulations of therapeutic DNA, has repeatedly demonstrated successful *in vitro* and *in vivo* gene transfection. While oral gene therapy has shown immense promise as treatment options in a variety of diseases, there are still significant barriers to overcome before it can be considered for clinical applications. In this review we provide an overview of the physiologic challenges facing the use of chitosan DNA nanoparticles for oral gene delivery at both the extracellular and intracellular level. From administration at the oral cavity, chitosan nanoparticles must traverse the gastrointestinal tract and protect its DNA contents from significant jumps in pH levels, various intestinal digestive enzymes, thick mucus layers with high turnover, and a proteinaceous glycocalyx meshwork. Once these extracellular barriers are overcome, chitosan DNA nanoparticles must enter intestinal cells, escape endolysosomes, and disassociate from genetic material at the appropriate time allowing transport of genetic material into the nucleus to deliver a therapeutic effect. The properties of chitosan nanoparticles and modified nanoparticles are discussed in this review. An understanding of the barriers to oral gene delivery and how to overcome them would be invaluable for future gene therapy development.

Key words: Gene therapy; Oral gene delivery; Chitosan nanoparticles; Glycocalyx; pH; Enzymes; Endolysosomal escape; Intracellular transport; Nuclear transport; Mucus

© The Author(s) 2016. Published by Baishideng Publishing

Group Inc. All rights reserved.

Core tip: Gene therapy is facing considerable challenges to realize its promise, the most important which is arguably achieving effective delivery of the therapeutic gene to the target cells. Therefore, developing alternative delivery strategies is crucial. This manuscript reviews the oral administration of chitosan DNA nanoparticles as a novel gene therapy delivery strategy.

Patel BJ, Vignesh NK, Hortelano G. Chitosan DNA nanoparticles for oral gene delivery. *World J Med Genet* 2016; 6(3): 22-33 Available from: URL: <http://www.wjgnet.com/2220-3184/full/v6/i3/22.htm> DOI: <http://dx.doi.org/10.5496/wjmg.v6.i3.22>

INTRODUCTION

Gene therapy, broadly understood as the delivery of nucleic acids for therapeutic purposes, has therapeutic potential for a number of medical conditions. Since its first clinical trials in 1990, gene therapy research has made a number of advancements^[1]. Despite its tribulations there has been some success, such as the treatment of immunodeficiencies through retroviral and lentiviral transduction of stem cells^[2], X-linked adrenoleukodystrophy through lentiviral transduction^[3], hemophilia through Adeno-associated virus delivery^[4], as well as the use of transduced T cells as a treatment for leukemia^[5]. Due to low cell survival upon transplantation and logistics inherent to *ex vivo* cell therapy, nucleic acids (including DNA, antisense oligonucleotides, and RNA interference systems) are typically delivered using viral vectors because of the high expression they induce in target cells^[6]. However, the ability of viral vectors to transduce cells and induce transgene expression may also trigger immune responses, and the potential activation of cancer-promoting genes in the case of integrating viral vectors^[7]. The safety of current gene therapy delivery systems, particularly that of integrating vectors is still not fully studied^[8]. Non-viral gene delivery lags behind viral vectors in targeting efficacy and transgene expression, which can be enhanced using mechanical techniques (e.g., use of electroporation or ultrasound) or chemical carriers (e.g., use of lipid, peptides, or polymers that protect the genetic material)^[9]. The use of non-viral DNA formulations for oral gene delivery is an area of active research interest^[10].

Oral gene delivery *via* chemical carriers has the potential to treat a number of inherited and acquired diseases^[11]. The gastrointestinal (GI) tract's large surface area provides a massive number of cells that can be potentially targeted. Successful delivery of therapeutic material to the GI is not restricted to the treatment of intestinal diseases such as inflammatory bowel disease, intestinal cancers, and intestinal symptoms of cystic fibrosis, but could also be potentially used as a treatment for systemic diseases (such as hemophilia) using the

extensive capillary network for efficient distribution of the transgenes^[6]. Additionally, effective exogenous DNA targeting of the intestinal stem cells in the crypts of Lieberkuhn in a stable manner could lead to potential long-term therapeutic gene expression^[6].

Despite such therapeutic potential, oral gene delivery must first overcome formidable extracellular and intracellular barriers in the GI tract. These include stomach acidity, glycocalyx, mucus, cell-uptake, endolysosomal escape, and nuclear import. The coupling of nucleic acids to polymers may assist in some of these challenges. Chitosan, a deacetylated form of chitin (Figure 1), which is derived from crustacean, has been shown to be an appropriate chemical polymer for overcoming such barriers. In addition to biodegradable and biocompatible properties^[12], this $\alpha(1-4)2$ -amino 2-deoxy β -glycan can efficiently compact DNA and has mucoadhesive and absorptive properties, buffering capacity, and membrane perturbing properties that make it a promising vehicle for oral gene delivery^[13]. This review discusses the various extracellular and intracellular barriers of oral gene delivery and how they can be overcome using chitosan and chitosan-modified nanoparticles.

EXTRACELLULAR BARRIERS

Free DNA is quickly denatured by the acidic pH in the stomach and intestinal and cellular enzymes, so it would have no therapeutic effect^[6]. However, chitosan-coupled therapeutic DNA sequences formulated as chitosan nanoparticles are effectively protected from various physiological and cellular barriers.

Acidity

pH varies greatly in the gastrointestinal tract, ranging from the extremely acidic stomach (mean pH 1.7) to the slightly alkaline ileum (mean pH 7.5) (Figure 2A)^[14-16]. Chitosan nanoparticles have been shown to protect enclosed nucleic acids across a wide range of pH. In the harsh stomach acid^[17,18] increased electrostatic interactions between positively charged chitosan and negatively charged DNA creates a more stable nanoparticle (Figure 2B). It is chitosan's amino-group with a pKa of 6.5 (Figure 1B) that accounts for protonation of the polymer at pHs below 6.5, and the resulting binding affinity between cationic chitosan and anionic DNA^[19].

As nanoparticles transit towards the small intestine, pH increases drastically from 1.7 in the stomach to 6.1 and 6.6 in the duodenum and jejunum respectively^[14,15]. During this shift, chitosan's amino groups are deprotonated, decreasing chitosan's binding affinity for DNA^[19,20]. Nanoparticle stability is slightly reduced, achieving an effective balance between DNA association and dissociation. It is at pH 6.8 and 7.0 that transfection efficiency of chondrocytes peaked^[21]. Interestingly however, Roy *et al.*^[17] reported transfection of stomach tissue by chitosan nanoparticles.

As the pH continues to climb distally in the intestine,




Figure 1 Chitin and chitosan molecular structure. The molecular structure of chitin (A) differs from chitosan (B) through its deacetylation. The resulting amino group on chitosan has a pKa of 6.5^[12].




Figure 2 Effect of gastrointestinal pH on chitosan-DNA nanoparticles. A: pH varies drastically throughout the human gastrointestinal tract from acidic stomach to slightly basic distal bowel^[14,15]. This in turn influences how chitosan-DNA polymers behave intraluminally; B: The conceptual interaction between negatively charged DNA and positively charged chitosan that becomes protonated at acidic pHs. This allows chitosan-DNA nanoparticles to remain stable at lower pHs.

chitosan deprotonates further until it is neutral. Neutral chitosan nanoparticles are quite unstable as they easily dissociate from DNA and begin to aggregate with themselves, no longer inhibited by positive-positive repulsive forces (Figure 2B)^[21,22]. Transfection efficiency also decreases at these high pH levels as has been demonstrated in human-lung carcinoma A549 cells (transfection at pH 7.6 < transfection at pH 6.9)^[23] and HEK 293 cells (transfection at pH 7.6 < transfection at pH 6.5)^[20].

Therefore, chitosan (pKa 6.5) is able to withstand a wide pH range from 1.5 (highly stabilized) to 6.8 (intermediate stability - optimal for transfection), before becoming destabilized and aggregating. In support of this model, *ex vivo* experiments on gastrointestinal tissue internalization of chitosan nanoparticles found internalization to be higher for the jejunum (pH 6.6) than the ileum (pH 7.5 - too basic) and duodenum (pH 6.1 - too acidic)^[24].

Mucus

The gastrointestinal mucus, composed of mucin fibers linked to various proteoglycans^[25], is constantly

produced, secreted and shed. This rapid turnover combined with the thickness of the mucus layer (greatest in the colon at 830 μ m) must be overcome by chitosan nanoparticles for effective oral gene delivery^[26]. There are three main mechanisms that have been explored.

For a long time mucoadhesion was thought to improve absorption by extending nanoparticle residence time in the GI tract and increasing bioavailability of nanoparticles for passive paracellular uptake^[12,27]. Therefore, researchers sought to improve mucoadhesion by formulating nanoparticles capable of greater interactions with mucus through electrostatic bonds, hydrogen bonds, hydrophobic interactions, and van der Waals forces^[28]. Chitosan is reported to have strong mucoadhesive properties^[27]. However mucoadhesion, on its own, will not solve this barrier to oral gene delivery, and may in fact impede drug delivery. This is because despite optimal mucoadhesion, nanoparticle delivery is limited by the rapid turnover time of mucus (50 to 270 min)^[29] during which the luminal layer of mucus is sloughed off taking the adherent nanoparticles with it (Figure 3)^[27].

Mucopenetration, or the ability to traverse the mucus




Figure 3 Gastrointestinal mucus as a barrier to chitosan-DNA nanoparticles transfection. Overcoming the mucus layer for oral gene delivery can be accomplished through three key mechanisms. Mucoadhesive nanoparticles adhere to mucus through chemical interactions. Bioavailability is limited by time, because intestinal mucus has a rapid turnover and adhered nanoparticles would be sloughed off with the luminal layer of mucus^[27]. Mucopenetration involves nanoparticles that are able to traverse intestinal mucus due to chemical properties. These nanoparticles do improve bioavailability even with time^[27]. Mucodisruptive nanoparticles cross the mucus layer by disrupting mucus structure. This can be accomplished by various mucolytic molecules and enzymes. Bioavailability increases with this mechanism, however intestinal mucus may be affected in the long-term^[27].

layer (Figure 3), has become the focus of overcoming the mucosal barrier in the last few years. Mucopenetration is achieved by creating a “slippery” nanoparticle that, in contrast to mucoadhesion, has decreased interactions with mucus^[30]. Chitosan was combined with hydrophilic 5 kDa polyethylene glycol (PEG) which would have decreased interactions with hydrophobic mucus and would not be sterically hindered^[30]. Nanoparticles with greater PEGylation permeated fresh mucus 5-fold compared with non-PEGylated nanoparticles (Figure 4A)^[30]. Pereira de Sousa *et al.*^[31] combined chitosan with chondroitin sulphate to create a densely charged but overall neutral nanoparticle surface that would theoretically have minimal

interactions with mucus. Chitosan chondroitin sulphate nanoparticles permeated to a greater depth (4-6 mm) of fresh undiluted porcine intestinal mucus compared with PDLG “purasorb” chitosan nanoparticles (Figure 4B)^[31]. Importantly, mucopenetration research remains in its early stages and further studies are required. To our knowledge no studies evaluating the effect of pDNA on the performance of chitosan nanoparticles modulated for mucopenetration have been conducted. This would be important to assess in the future.

Finally disruption of the mucosal layer can be achieved by combining nanoparticles with mucolytic agents (Figure 3). Studies have shown enhanced mucodisruption ability




Figure 4 Effect of gastrointestinal enzymes on chitosan-DNA nanoparticles. A: Various digestive enzymes are secreted throughout the gastrointestinal tract. The genetic material within chitosan nanoparticles is most threatened by pancreatic nucleases such as deoxyribonuclease and ribonuclease^[34]; B: Conjugation of chitosan polymer with ethylenediaminetetraacetic acid through a covalent bond at the amino group^[37].

of nanoparticles when combined with mucolytic molecules (N-acetylcystein, L-cysteine, sulfhydryl compounds)^[32] and mucolytic enzymes (papain, bromelain, trypsin)^[33]. These studies, however were not conducted with chitosan. With the mucodisruption technique, reversibility of mucolysis must continue to be assessed in order to ensure that the intestinal mucus is not completely damaged and that it retains its protective function against viruses and pathogens^[32].

Enzymes

Administered genes are exposed to the harsh environment of the GI lumen, filled with resident degrading enzymes, those secreted into the lumen and those membrane bound^[34]. Unavoidably, therapeutic nanoparticles are also subject to the degradation processes of these enzymes and thus, pose a barrier that must be dealt with to improve the efficacy of oral gene delivery^[34].

Conjugation of enzyme inhibitors with carrier vehicles was proposed a long time ago^[35] and has been explored using a variety of enzyme inhibitors^[36]. These inhibitor conjugated chitosan nanoparticles have been shown to protect a variety of drug molecules (including pDNA^[34,37], insulin^[38-40], and calcitonin^[41-44]) and thereby promote therapeutic effect (Table 1). For oral gene delivery, inhibiting nuclease destruction of genetic material is most important and can be accomplished using the following inhibitors (ranked by efficacy): Ethylenediaminetetraacetic acid (EDTA) > sodium dodecyl sulphate (SDS) > aurotricarboxylic acid (ATA) > and poly(acrylic acid) (PAA)^[37]. Loretz *et al*^[37] demonstrated that 50% of pDNA was protected from nuclease destruction when complexed with EDTA-conjugated chitosan. This is compared to

almost complete destruction of unprotected pDNA^[37]. EDTA-chitosan nanoparticles also have 35% improved transfection efficiency in Caco-2 cells compared with unmodified chitosan nanoparticles^[45]. Similarly, Martien *et al*^[34] demonstrated almost complete protection of pDNA from lysozyme and deoxyribonuclease when packaged in ATA-chitosan conjugated nanoparticles compared with unmodified chitosan nanoparticles (complete destruction with both lysozyme and deoxyribonuclease). Conjugation of enzyme inhibitors to chitosan involves the formation of a covalent bond between a carboxylic acid group on the inhibitor and free amino acids on chitosan. Notably, the covalent conjugation ensures that the inhibitor is not orally absorbed^[37] and reduces the toxicity and side effects. This is supported by LDH-release cytotoxicity tests on Caco-2 cells that showed significantly less toxicity (< 1% over 4 h) for chitosan-EDTA particles and polymers compared with unmodified chitosan particles and polymers^[45]. It is important to note that prolonged administration of enzyme inhibitors has severe side effects including pancreatic hyperplasia and carcinoma and thus chronic effects of inhibitor-conjugated chitosan nanoparticles must be explored^[46].

Glycocalyx

The fibrous brush border glycocalyx (FBBG) is a 400-500 nm thick glycoprotein network that covers the apical surface of intestinal epithelial cells, particularly enterocytes (Figure 5). As visualised through high resolution scanning electron microscopy, the FBBG consists of mucin-like glycoproteins^[47] that originate from the tips of tightly packed microvilli and anastomose with neighbouring glycoproteins^[48]. The resulting intricate mesh has the

Table 1 Chitosan-enzyme inhibitor conjugates¹

	Enzyme inhibition	Drug protection
Chitosan-ATA	DNAase	pDNA ^[54]
	Lysozyme	
Chitosan-EDTA	DNAase	pDNA ^[57]
	Trypsin	Insulin ^[39]
	Chymotrypsin	Trypsin and bromelain ^[93]
Chitosan-aprotinin	Trypsin	Insulin ^[38]
	Chymotrypsin	Calcitonin ^[41]
Chitosan-BBI	Trypsin	Calcitonin ^[42]
	Chymotrypsin	Insulin ^[40]
Chitosan-pepstatin	Pepsin	Calcitonin ^[43]
Chitosan-elastinal	Elastase	Calcitonin ^[42]
		Insulin ^[40]
Chitosan-phenylboronic acid	Trypsin	Calcitonin ^[44]
	Elastase	
Chitosan-bacitracin		Hepatitis B surface antigen ^[94]

¹Summary of the various enzyme inhibitors that chitosan has been conjugated with in order to decrease degradation and maximize delivery of drug product. All references of original studies are included. Chitosan-ATA and chitosan-EDTA conjugates are most relevant to gene therapy advancements, as they have been shown to protect pDNA and increase transfection efficiency. EDTA: Ethylenediaminetetraacetic acid; ATA: Aurintricarboxylic acid; BBI: Bowman-birk inhibitor.

potential to affect transfection efficiency of chitosan nanoparticles. The mechanism by which this would occur is unclear as existing data is mixed.

Originally, the glycocalyx was theorized to act as a size-selective diffusion barrier, with pores between 7.4 and 28.8 nm wide, allowing only appropriately sized particles to pass through and access transmembrane receptors located on microvilli^[49]. More recent studies however have demonstrated that the glycocalyx does not have a diffusion effect on drug delivery (in particular insulin) but rather has a digestive effect on the therapeutic molecule due increased concentration of digestive enzymes at the FBBG^[50]. More research is required to better understand this potential barrier to oral gene delivery.

The data on how to overcome the intestinal glycocalyx barrier for improved drug delivery is also mixed. Some studies promote the theory that increased interaction with glycocalyx layer, similar to that of bacteria and viruses, promotes effective drug delivery due to increased bioavailability of drugs at the apical membrane^[49]. In contrast, other research demonstrates that electrostatic interactions between anionic glycocalyx and cationic nanoparticles would decrease nanoparticle charge, stimulate premature release of DNA, and cause DNA relaxation^[51]. Overall, more research is required on the glycocalyx and its impact on transfection efficiency of nanoparticles.

INTRACELLULAR BARRIERS

Transport and cellular uptake

The process of cellular uptake of the nanoparticle is critical, such that its genetic component is internalized

and functionally available to elicit the desired biological effect. In addition to the key effect of pH on transfection^[52] nanoparticles are internalized *via* a receptor-mediated endocytosis^[53,54]. However, it is important to note that nanoparticles of different formulations - and sizes - may not necessarily share the same mechanism for cell entry^[55]. Thus, chitosan may have the ability to interact with the cell membrane *via* electrostatic forces leading to an energy dependent adsorptive endocytosis process. Additionally, Nam *et al.*^[55] showed that the uptake process of chitosan nanoparticles involved multiple distinct pathways, including the clathrin-mediated endocytosis, macropinocytosis and caveolae-mediated endocytosis. Further, they showed that each pathway was independent from the other.

In addition to delivering therapeutic nucleic acids to intestinal cells, ingested chitosan nanoparticles may also reach more distal sites by crossing the intestinal epithelium and entering the circulation system. This may be achieved through paracellular and/or transcellular transport of nanoparticles^[56].

Paracellular transport is improved by chitosan's permeation enhancing properties. Positively charged chitosan interacts with tight junction proteins (occludin, ZO-1, and F-actin) and destabilizes plasma membranes to decrease the trans-epithelial electrical resistance (TEER) of cell monolayers^[54]. Increased paracellular transport of macromolecules such as insulin^[57] has been shown to consistently occur when chitosan is in solution. The permeability influence of chitosan-nanoparticles is less conclusive, but a study by Vllasaliu *et al.*^[58] showed that both chitosan and chitosan nanoparticles are effective at opening tight-junctions. Chitosan permeability depends on electrostatic interactions between chitosan and tight junction proteins, and thus, the pore-opening process may be affected by the degree of deacetylation and surface area of charge exposed on chitosan nanoparticles^[58]. Further, nanoparticle size may inhibit paracellular transport due to maximal widening of tight-junction pores^[58]. Thus far, few safety issues have been identified with chitosan's ability to traverse across the intestinal epithelium. This is because chitosan's ability to open tight junctions is reversible^[58], and because common negatively charged GI endotoxins (such as lipopolysaccharide) are unable to cross the intestinal epithelium along with chitosan nanoparticles^[59].

Transcellular transport may occur through transcytosis^[60], which involves cellular uptake (pinocytosis, clathrin-mediated endocytosis, discussed earlier) at the apical membrane, transport across the cell interior, and release at the basolateral pole^[56]. Although M-cells make up only 1% of the total intestinal surface, it is believed that most transcytosis occurs with these cells due to higher rates of endocytosis^[56]. Transcytosis efficiency decreases with nanoparticle size^[56] and increases with nanoparticle stability^[61].

Endolysosome escape

DNA escape from the endolysosomal pathway is a key barrier to chitosan gene delivery. It is theorized that




Figure 5 Glycocalyx effect on nanoparticles.

A thick mesh like structure of glycocalyx proteins lines the apical membrane of the small bowel with pores 7-29 nm in size^[49]. It is unclear whether the glycocalyx has a size effect on diffusion, as existing literature contains mixed results.

chitosan nanoparticles may escape from endosomes through a "Proton Sponge Effect"^[62,63]. According to this theory, endocytosed chitosan nanoparticles become protonated upon fusion with acidic lysosomes. This is followed by an influx of cytosolic H^+ , Cl^- and H_2O , to maintain the endosomal pH, charge, and osmolarity, respectively^[64]. Such influx causes the lysosome to swell and rupture, thereby releasing nanoparticles into the cell cytosol (Figure 6)^[64]. Although the proton sponge effect is supported by cationic polymers with strong buffering capacities (e.g., PEI) chitosan has a relatively low buffering capacity over a narrow pH range (pH 5-7)^[65]. This makes it a poor "proton-sponge" at endosomal pH, only having a limited influx of protons^[65]. This is reflected in its delayed endosomal escape and transfection, observed in HEK 293 cells treated with chitosan and PEI^[65]. Chitosan endosomal rupture occurred at 72 h-post-treatment, compared with PEI's escape within 24 h post-treatment^[65], a delay perhaps required for a sufficient amount of chitosan to accumulate to induce a proton-sponge rupture.

Proton sponge endosomal escape can be enhanced through the addition of high buffering-capacity groups. For example, studies have examined the effect of the biocompatible and buffering imidazole ring (pKa 6.15) on the release of chitosan nanoparticles into the cytoplasm^[66]. In separate studies, increased expression was observed for chitosan-graft-histadine^[67] and chitosan-graft-urocanic acid^[66], both of which bear the imidazole ring. Although expression was improved relative to chitosan, improvement was cell-specific and neither complex was able to reach the expression-levels of lipofectamine^[67,68]. Confocal microscopy and treatment with bafilomycin A (proton pump inhibitor) confirmed that histadine's effect occurs due to improved endosomal escape of complexes^[67]. Multiple experiments have also been conducted on PEI-chitosan hybrids including chitosan/PEI mixes and chitosan-graft-PEI systems. PEI is an effective non-viral vector due to its high buffering capacity (multiple amino groups) and enhanced endosomal escape properties^[68]. However, PEI's effect is limited *in vivo*, due to its cytotoxicity in multiple cell

lines^[69]. Thus, combining PEI with chitosan, known for its biocompatibility and biodegradability, has shown to be particularly effective with synergistic results^[68]. Compared with PEI, chitosan-graft-PEI has higher transfection efficiency *in vitro* and *in vivo*^[70], and has decreased cytotoxicity across three cell lines (293 T, HeLa and HepG2)^[71]. Transfection efficiency even reached that of lipofectamine in the human HEK293 cell line^[71].

In addition to the described proton-sponge effect, there are other means by which nanoparticles escape from endosomes. Membrane destabilization by chitosan is supported by experiments tracking endocytosed nanoparticles^[72,73]. In these experiments, some DNA is released from early endosomes^[72] before the proton-sponge effect can be triggered by acidic lysosomes. Modifications to improve chitosan's membrane perturbant properties include addition of chloroquine^[74], poly(propylacrylic acid)^[75], and synthetic peptides^[76]. Synthetic peptides are pH sensitive, changing conformation from random coils (at physiological pH) to alpha helices (at endosomal acidity) that interact with and destabilize the membrane through pore-formation^[77]. Second, free chitosan polymer may also aid endosomal escape. Unlike protonated nanoparticles (stabilized by DNA) that decrease in volume, free chitosan expands from its coiled form due to repulsive forces^[78]. This directly increases osmotic pressure and leads to endosomal rupture^[78]. Thibault *et al.*^[79] added free chitosan at different time-points to HEK 293 cells already treated with inefficient chitosan nanoparticles (< 5 N:P levels)^[79]. Addition of free chitosan at 4 and 8 h post-nanoparticle treatment resulted in a near 100% rescue of transfection, reaching similar levels to the optimal N:P of 5^[79].

DNA-release

In addition to the protective role of chitosan discussed above it is also important for chitosan to have a timely release of DNA. Nanoparticles of intermediate stability, associated with intermediate molecular weight and degree of deacytylation appropriately release DNA in synchrony with lysosomal escape and achieve efficient transfection^[80]. This is in contrast to the inefficient transfection ability of




Figure 6 Proton sponge effect. Conceptual image of how chitosan-DNA nanoparticles may escape from lysosomes once endocytosed. H⁺, Cl⁻, and H₂O influx into the lysosome in order to maintain pH, charge, and osmolarity. This movement of molecules causes swelling of the lysosome, eventual bursting, and escape from lysosomal digestion^[77].

hyperstabilised and hypostabilized nanoparticles. Up to 80% of hyperstabilized nanoparticles (high molecular weight and degree of deacetylation) remained associated after lysosomal escape, and were too large to enter the nucleus^[80]. Hypostabilized nanoparticles (low molecular weight and/or degree of deacetylation) dissociated prematurely before endolysosomal escape and were degraded^[80]. Thus, chitosan nanoparticles of intermediate stability are optimal for timely release of DNA and efficient transfection.

Nuclear import

In order to have a therapeutic effect, DNA must then enter the nucleus and be accessible to transcriptional enzymes, while siRNA - but not DNA coding for siRNA - exerts its inhibitory function in the cytoplasm. Considering that the majority of GI cells (*e.g.*, enterocytes, goblet cells, paneth cells) are post-mitotic, genetic material must pass through nuclear pores, as opposed to waiting for the nuclear envelope to disintegrate during cell division. Import of material (> 40 kDa) through the nuclear pore occurs through indirect active transport^[81]. It involves interactions between importins and proteins containing nuclear localization signal (NLS) (Figure 7A).

Therefore, enhanced nuclear import of pDNA is an important goal. Sequences where NLS-containing proteins bind are often added to therapeutic pDNA. The 72 bp SV40 enhancer sequence upon which many mammalian transcription factors (AP1, AP2, NF- κ B, Oct1, TEF1, each containing NLS sequences) bind has been used to enhance the import of perinuclear localized pDNA into the nucleus through interaction with importin proteins^[80,82]. Opanasopit *et al.*^[83] showed increased

transfection efficiency of nanoparticles containing DNA and NLS-containing peptides. Additionally, it is possible to enhance gene expression by covalently complex NLS-containing proteins onto the pDNA strand itself^[83]. However, it is important for the protein's interaction with pDNA to be spatially distinct from its NLS-binding sequence so that importins are not sterically hindered from binding (Figure 7B).

Interestingly, some evidence exists for nuclear import of entire chitosan/DNA complexes. Zhao *et al.*^[84] found increased localization of complexes when a synthetic protein, containing SV40 NLS signal, was covalently linked to chitosan. Further addition of kinases promoted intra-nuclear disassociation and improved transfection efficacy beyond that of lipofectamine^[84]. Chitosan localized in the nucleus did not interfere with gene expression^[73].

Immune response

Chitosan is a biodegradable polysaccharide that has been administered to individuals without any adverse side effects^[85]. Chitosan is also naturally degraded through lysosome digestion according to its molecular weight and degree of deacetylation. These characteristics of chitosan have triggered the utilization of chitosan as an adjuvant in vaccines^[86]. The viscous nature of chitosan promotes the formation of an antigen depot. Injection of antigen with chitosan resulted in 60% of the antigen remaining at the injection site for 7 d whereas less than 9% of the antigen remained after 8 h when it was administered with saline^[86]. In combination with antigens, chitosan has been demonstrated to induce activation of NK cells, macrophages, inducing cytokines, stimulating antibody




Figure 7 Model of DNA combined with importin/nuclear localization signal complex to enhance importation of DNA through nucleopores. A: Plasmid DNA bound to importins/nuclear localization signal (NLS) complex through transcription factors^[79,81]; B: Plasmid DNA bound to importins/NLS complex through a synthetic protein^[83].

production and cytotoxic T lymphocyte responses^[87]. Chitosan has been shown to elicit a combination TH1/TH2 response, thus promoting both humoral and cell mediated immunity. The increased immune response has been reported to produce a high titer antibody response to the antigen^[88].

When dealing with oral vaccination, the uptake by the Peyer's patch is a crucial step in stimulation an immune response. Studies have developed chitosan particles that have the optimal size and zeta potential to be targeted to the M-cells of the Peyer's patch^[89]. Particles that are < 10 μm are taken up by M-cells while those < 5 μm can be transported to the spleen or local lymph nodes to stimulate IgM and IgG production^[89]. As a result, chitosan promises to serve as a carrier system of antigens and/or antigen-coding genetic material to act as an adjuvant in vaccinations to enhance both cell mediated as well as humoral immune responses. However, chitosan's immunomodulatory effects have also been shown in studies that utilize the nasal administration of chitosan to induce the immune system^[90]. Chitosan is also a viable vehicle to be used in oral administration due to its adhesive and transport properties^[17]. Its stability when complexed with DNA make it suitable for the delivery of the cargo to the M-cells of the GI-tract, allowing for the generation of both mucosal as well as systemic immunity^[17]. The immunity of chitosan nanoparticles has also been exploited to induce oral tolerance. Goldmann *et al.*^[91] fed mice with oral nanoparticles complexed to ovalbumin (OVA) DNA. Despite OVA being a very good antigen, the nanoparticle treatment suppressed a humoral response after subsequent exposure to OVA protein. The authors noticed a shift from TH₁ to a TH₂/TH₃ together with the induction of OVA-specific regulatory T cells^[91]. The same group took this work a step further by protecting mice against abdominal aortic transplantation after feeding mice with chitosan nanoparticles containing DNA coding for k(b), one of the MHC-I molecules of the

donor^[92]. These results open new applications for oral chitosan nanoparticles.

CONCLUSION

Oral delivery of genes specifically is an intriguing technique to treat GI pathologies as well as other chronic systemic diseases such as haemophilia^[12,18]. However, it should be noted that hormone related disorders are much more challenging to treat due to the strict regulation the protein expression and the systemic affect that many hormones exhibit. While oral gene therapy has shown immense promise as treatment options in a variety of diseases, there are still significant barriers to overcome before it can be considered for clinical applications. By modulating the physical and chemical characteristics of the nanoparticle both extracellular and intracellular barriers may be circumvented. By judicious design of the nanoparticles therapeutic potential for the targeting of many systemic diseases may be possible. Further research in this field is required in order to develop in the future chitosan nanoparticles of viable clinical use.

ACKNOWLEDGMENTS

The authors thank the Canadian Hemophilia Society Bayer Inc. and Nazarbayev University ORAU for their supporting to this work.

REFERENCES

- 1 **Herzog RW**, Cao O, Srivastava A. Two decades of clinical gene therapy--success is finally mounting. *Discov Med* 2010; **9**: 105-111 [PMID: 20193635]
- 2 **Fischer A**, Hacein-Bey-Abina S, Cavazzana-Calvo M. Gene therapy of primary T cell immunodeficiencies. *Gene* 2013; **525**: 170-173 [PMID: 23583799 DOI: 10.1016/j.gene.2013.03.092]
- 3 **Cartier N**, Hacein-Bey-Abina S, Bartholomae CC, Veres G, Schmidt

- M, Kutschera I, Vidaud M, Abel U, Dal-Cortivo L, Caccavelli L, Mahlaoui N, Kiermer V, Mittelstaedt D, Bellesme C, Lahlou N, Lefrère F, Blanche S, Audit M, Payen E, Leboulch P, l'Homme B, Bougnères P, Von Kalle C, Fischer A, Cavazzana-Calvo M, Aubourg P. Hematopoietic stem cell gene therapy with a lentiviral vector in X-linked adrenoleukodystrophy. *Science* 2009; **326**: 818-823 [PMID: 19892975 DOI: 10.1126/science.1171242]
- 4 **Nathwani AC**, Tuddenham EG, Rangarajan S, Rosales C, McIntosh J, Linch DC, Chowdary P, Riddell A, Pie AJ, Harrington C, O'Beirne J, Smith K, Pasi J, Glader B, Rustagi P, Ng CY, Kay MA, Zhou J, Spence Y, Morton CL, Allay J, Coleman J, Sleep S, Cunningham JM, Srivastava D, Basner-Tschakarjan E, Mingozzi F, High KA, Gray JT, Reiss UM, Nienhuis AW, Davidoff AM. Adenovirus-associated virus vector-mediated gene transfer in hemophilia B. *N Engl J Med* 2011; **365**: 2357-2365 [PMID: 22149959 DOI: 10.1056/NEJMoa1108046]
- 5 **Ramos CA**, Savoldo B, Dotti G. CD19-CAR trials. *Cancer J* 2004; **20**: 112-118 [PMID: 24667955 DOI: 10.1097/PPO.000000000000031]
- 6 **O'Neill MJ**, Bourre L, Melgar S, O'Driscoll CM. Intestinal delivery of non-viral gene therapeutics: physiological barriers and preclinical models. *Drug Discov Today* 2011; **16**: 203-218 [PMID: 21262379 DOI: 10.1016/j.drudis.2011.01.003]
- 7 **Cavazzana-Calvo M**, Thrasher A, Mavilio F. The future of gene therapy. *Nature* 2004; **427**: 779-781 [PMID: 14985734 DOI: 10.1038/427779a]
- 8 **Rothe M**, Schambach A, Biasco L. Safety of gene therapy: new insights to a puzzling case. *Curr Gene Ther* 2014; **14**: 429-436 [PMID: 25245088 DOI: 10.2174/1566523214666140918110905]
- 9 **Niïdome T**, Huang L. Gene therapy progress and prospects: nonviral vectors. *Gene Ther* 2002; **9**: 1647-1652 [PMID: 12457277 DOI: 10.1038/sj.gt.3301923]
- 10 **Junquera E**, Aicart E. Recent progress in gene therapy to deliver nucleic acids with multivalent cationic vectors. *Adv Colloid Interface Sci* 2016; **233**: 161-175 [PMID: 26265376 DOI: 10.1016/j.cis.2015.07.003]
- 11 **Dolgin E**. Immunology: Oral solutions. *Nature* 2014; **515**: S166-S167 [PMID: 25427208 DOI: 10.1038/515S166a]
- 12 **Hejazi R**, Amiji M. Chitosan-based gastrointestinal delivery systems. *J Control Release* 2003; **89**: 151-165 [PMID: 12711440 DOI: 10.1016/S0168-3659(03)00126-3]
- 13 **Mao S**, Sun W, Kissel T. Chitosan-based formulations for delivery of DNA and siRNA. *Adv Drug Deliv Rev* 2010; **62**: 12-27 [PMID: 19796660 DOI: 10.1016/j.addr.2009.08.004]
- 14 **Dressman JB**, Berardi RR, Dermentzoglou LC, Russell TL, Schmaltz SP, Barnett JL, Jarvenpaa KM. Upper gastrointestinal (GI) pH in young, healthy men and women. *Pharm Res* 1990; **7**: 756-761 [PMID: 2395805 DOI: 10.1023/A: 1015827908309]
- 15 **Evans DF**, Pye G, Bramley R, Clark AG, Dyson TJ, Hardcastle JD. Measurement of gastrointestinal pH profiles in normal ambulant human subjects. *Gut* 1988; **29**: 1035-1041 [PMID: 3410329 DOI: 10.1136/gut.29.8.1035]
- 16 **Kong F**, Singh RP. Disintegration of solid foods in human stomach. *J Food Sci* 2008; **73**: R67-R80 [PMID: 18577009 DOI: 10.1111/j.1750-3841.2008.00766.x]
- 17 **Roy K**, Mao HQ, Huang SK, Leong KW. Oral gene delivery with chitosan-DNA nanoparticles generates immunologic protection in a murine model of peanut allergy. *Nat Med* 1999; **5**: 387-391 [PMID: 10202926 DOI: 10.1038/7385]
- 18 **Dhadwar SS**, Kiernan J, Wen J, Hortelano G. Repeated oral administration of chitosan/DNA nanoparticles delivers functional FVIII with the absence of antibodies in hemophilia A mice. *J Thromb Haemost* 2010; **8**: 2743-2750 [PMID: 20961391 DOI: 10.1111/j.1538-7836.2010.04116.x]
- 19 **Köping-Höggård M**, Mel'nikova YS, Vårum KM, Lindman B, Artursson P. Relationship between the physical shape and the efficiency of oligomeric chitosan as a gene delivery system in vitro and in vivo. *J Gene Med* 2003; **5**: 130-141 [PMID: 12539151 DOI: 10.1002/jgm.327]
- 20 **Lavertu M**, Méthot S, Tran-Khanh N, Buschmann MD. High efficiency gene transfer using chitosan/DNA nanoparticles with specific combinations of molecular weight and degree of deacetylation. *Biomaterials* 2006; **27**: 4815-4824 [PMID: 16725196 DOI: 10.1016/j.biomaterials.2006.04.029]
- 21 **Zhao X**, Yu SB, Wu FL, Mao ZB, Yu CL. Transfection of primary chondrocytes using chitosan-pEGFP nanoparticles. *J Control Release* 2006; **112**: 223-228 [PMID: 16556468 DOI: 10.1016/j.jconrel.2006.01.016]
- 22 **Strand SP**, Danielsen S, Christensen BE, Vårum KM. Influence of chitosan structure on the formation and stability of DNA-chitosan polyelectrolyte complexes. *Biomacromolecules* 2005; **6**: 3357-3366 [PMID: 16283766 DOI: 10.1021/bm0503726]
- 23 **Sato T**, Ishii T, Okahata Y. In vitro gene delivery mediated by chitosan. effect of pH, serum, and molecular mass of chitosan on the transfection efficiency. *Biomaterials* 2001; **22**: 2075-2080 [PMID: 11432586 DOI: 10.1016/S0142-9612(00)00385-9]
- 24 **Behrens I**, Pena AI, Alonso MJ, Kissel T. Comparative uptake studies of bioadhesive and non-bioadhesive nanoparticles in human intestinal cell lines and rats: the effect of mucus on particle adsorption and transport. *Pharm Res* 2002; **19**: 1185-1193 [PMID: 12240945 DOI: 10.1023/A: 1019854327540]
- 25 **Bowman K**, Leong KW. Chitosan nanoparticles for oral drug and gene delivery. *Int J Nanomedicine* 2006; **1**: 117-128 [PMID: 17722528 DOI: 10.2147/nano.2006.1.2.117]
- 26 **Atuma C**, Strugala V, Allen A, Holm L. The adherent gastrointestinal mucus gel layer: thickness and physical state in vivo. *Am J Physiol Gastrointest Liver Physiol* 2001; **280**: G922-G929 [PMID: 11292601]
- 27 **Lai SK**, Wang YY, Hanes J. Mucus-penetrating nanoparticles for drug and gene delivery to mucosal tissues. *Adv Drug Deliv Rev* 2009; **61**: 158-171 [PMID: 19133304 DOI: 10.1016/j.addr.2008.11.002]
- 28 **Lai SK**, O'Hanlon DE, Harrold S, Man ST, Wang YY, Cone R, Hanes J. Rapid transport of large polymeric nanoparticles in fresh undiluted human mucus. *Proc Natl Acad Sci USA* 2007; **104**: 1482-1487 [PMID: 17244708 DOI: 10.1073/pnas.0608611104]
- 29 **Lehr CM**, Poelma FGJ, Junginger HE, Tukker JJ. An estimate of turnover time of intestinal mucus gel layer in the rat in situ loop. *Int J Pharm* 1991; **70**: 235-240 [DOI: 10.1016/0378-5173(91)90287-X]
- 30 **Pereira de Sousa I**, Moser T, Steiner C, Fichtl B, Bernkop-Schnürch A. Insulin loaded mucus permeating nanoparticles: Addressing the surface characteristics as feature to improve mucus permeation. *Int J Pharm* 2016; **500**: 236-244 [PMID: 26802494 DOI: 10.1016/j.ijpharm.2016.01.022]
- 31 **Pereira de Sousa I**, Steiner C, Schmutzler M, Wilcox MD, Veldhuis GJ, Pearson JP, Huck CW, Salvenmoser W, Bernkop-Schnürch A. Mucus permeating carriers: formulation and characterization of highly densely charged nanoparticles. *Eur J Pharm Biopharm* 2015; **97**: 273-279 [PMID: 25576256 DOI: 10.1016/j.ejpb.2014.12.024]
- 32 **Sheffner AL**. The reduction in vitro in viscosity of mucoprotein solutions by a new mucolytic agent, N-acetyl-L-cysteine. *Ann N Y Acad Sci* 1963; **106**: 298-310 [PMID: 13977050]
- 33 **Samaridou E**, Karidi K, de Sousa IP, Cattoz B, Griffiths P, Kammona O, Bernkop-Schnürch A, Kiparissides C. Enzyme functionalized PLGA nanoparticles with enhanced mucus permeation rate. *Nano LIFE* 2014; **4**: 1441013 [DOI: 10.1142/S179398441441013X]
- 34 **Martien R**, Loretz B, Schnürch AB. Oral gene delivery: design of polymeric carrier systems shielding toward intestinal enzymatic attack. *Biopolymers* 2006; **83**: 327-336 [PMID: 16609969 DOI: 10.1002/bip.20521]
- 35 **Bernkop-Schnürch A**, Paikl C, Valenta C. Novel bioadhesive chitosan-EDTA conjugate protects leucine enkephalin from degradation by aminopeptidase N. *Pharm Res* 1997; **14**: 917-922 [PMID: 9244150 DOI: 10.1023/A: 1012108118670]
- 36 **Werle M**, Takeuchi H, Bernkop-Schnürch A. Modified chitosans for oral drug delivery. *J Pharm Sci* 2009; **98**: 1643-1656 [PMID: 18781621 DOI: 10.1002/jps.21550]
- 37 **Loretz B**, Föger F, Werle M, Bernkop-Schnürch A. Oral gene delivery: Strategies to improve stability of pDNA towards intestinal digestion. *J Drug Target* 2006; **14**: 311-319 [PMID: 16882551 DOI: 10.1080/10611860600823766]
- 38 **Werle M**, Loretz B, Entstrasser D, Föger F. Design and evaluation of a chitosan-aprotinin conjugate for the peroral delivery of therapeutic peptides and proteins susceptible to enzymatic degradation. *J Drug*

- Target* 2007; **15**: 327-333 [PMID: 17541841 DOI: 10.1080/10611860701349141]
- 39 **Young LS**, Searle PF, Onion D, Mautner V. Viral gene therapy strategies: from basic science to clinical application. *J Pathol* 2006; **208**: 299-318 [PMID: 16362990 DOI: 10.1002/path.1896]
- 40 **Guggi D**, Kast CE, Bernkop-Schnürch A. In vivo evaluation of an oral salmon calcitonin-delivery system based on a thiolated chitosan carrier matrix. *Pharm Res* 2003; **20**: 1989-1994 [PMID: 14725364 DOI: 10.1023/B: PHAM.000008047.82334.7d]
- 41 **Bernkop-Schnürch A**, Giovannelli R, Valenta C. Peroral administration of enzymes: strategies to improve the galenic of dosage forms for trypsin and bromelain. *Drug Dev Ind Pharm* 2000; **26**: 115-121 [PMID: 10697748 DOI: 10.1081/DDC-100100336]
- 42 **Werle M**, Takeuchi H. Chitosan-aprotinin coated liposomes for oral peptide delivery: Development, characterisation and in vivo evaluation. *Int J Pharm* 2009; **370**: 26-32 [PMID: 19073243 DOI: 10.1016/j.ijpharm.2008.11.013]
- 43 **Krauland AH**, Guggi D, Bernkop-Schnürch A. Oral insulin delivery: the potential of thiolated chitosan-insulin tablets on non-diabetic rats. *J Control Release* 2004; **95**: 547-555 [PMID: 15023465 DOI: 10.1016/j.jconrel.2003.12.017]
- 44 **Guggi D**, Krauland AH, Bernkop-Schnürch A. Systemic peptide delivery via the stomach: in vivo evaluation of an oral dosage form for salmon calcitonin. *J Control Release* 2003; **92**: 125-135 [PMID: 14499191 DOI: 10.1016/S0168-3659(03)00299-2]
- 45 **Loretz B**, Bernkop-Schnürch A. In vitro evaluation of chitosan-EDTA conjugate polyplexes as a nanoparticulate gene delivery system. *AAPS J* 2006; **8**: E756-E764 [PMID: 17285741 DOI: 10.1208/aapsj080485]
- 46 **Melmed RN**, El-Aaser AA, Holt SJ. Hypertrophy and hyperplasia of the neonatal rat exocrine pancreas induced by orally administered soybean trypsin inhibitor. *Biochim Biophys Acta* 1976; **421**: 280-288 [PMID: 1252469 DOI: 10.1016/0304-4165(76)90294-4]
- 47 **Maury J**, Nicoletti C, Guzzo-Chambraud L, Maroux S. The filamentous brush border glycocalyx, a mucin-like marker of enterocyte hyper-polarization. *Eur J Biochem* 1995; **228**: 323-331 [PMID: 7535695 DOI: 10.1111/j.1432-1033.1995.tb20267.x]
- 48 **Horiuchi K**, Naito I, Nakano K, Nakatani S, Nishida K, Taguchi T, Ohtsuka A. Three-dimensional ultrastructure of the brush border glycocalyx in the mouse small intestine: a high resolution scanning electron microscopic study. *Arch Histol Cytol* 2005; **68**: 51-56 [PMID: 15827378 DOI: 10.1679/aohc.68.51]
- 49 **Frey A**, Giannasca KT, Weltzin R, Giannasca PJ, Reggio H, Lencer WI, Neutra MR. Role of the glycocalyx in regulating access of microparticles to apical plasma membranes of intestinal epithelial cells: implications for microbial attachment and oral vaccine targeting. *J Exp Med* 1996; **184**: 1045-1059 [PMID: 9064322]
- 50 **Aoki Y**, Morishita M, Takayama K. Role of the mucous/glycocalyx layers in insulin permeation across the rat ileal membrane. *Int J Pharm* 2005; **297**: 98-109 [PMID: 15907598 DOI: 10.1016/j.ijpharm.2005.03.004]
- 51 **Calderon AJ**, Baig M, Pichette B, Muzykantov V, Muro S, Eckmann DM. Effect of Glycocalyx on Drug Delivery Carriers Targeted to Endothelial Cells. *Int J Transp Phenom* 2011; **12**: 63-75 [PMID: 22679359]
- 52 **Nimesh S**, Thibault MM, Lavertu M, Buschmann MD. Enhanced gene delivery mediated by low molecular weight chitosan/DNA complexes: effect of pH and serum. *Mol Biotechnol* 2010; **46**: 182-196 [PMID: 20454872 DOI: 10.1007/s12033-010-9286-1]
- 53 **Ishii T**, Okahata Y, Sato T. Mechanism of cell transfection with plasmid/chitosan complexes. *Biochim Biophys Acta* 2001; **1514**: 51-64 [PMID: 11513804 DOI: 10.1016/S0005-2736(01)00362-5]
- 54 **Oh JM**, Choi SJ, Kim ST, Choy JH. Cellular uptake mechanism of an inorganic nanovehicle and its drug conjugates: Enhanced efficacy due to clathrin-mediated endocytosis. *Bioconjug Chem* 2006; **17**: 1411-1417 [PMID: 17105218 DOI: 10.1021/bc0601323]
- 55 **Nam HY**, Kwon SM, Chung H, Lee SY, Kwon SH, Jeon H, Kim Y, Park JH, Kim J, Her S, Oh YK, Kwon IC, Kim K, Jeong SY. Cellular uptake mechanism and intracellular fate of hydrophobically modified glycol chitosan nanoparticles. *J Control Release* 2009; **135**: 259-267 [PMID: 19331853 DOI: 10.1016/j.jconrel.2009.01.018]
- 56 **des Rieux A**, Fievez V, Garinot M, Schneider YJ, Préat V. Nanoparticles as potential oral delivery systems of proteins and vaccines: a mechanistic approach. *J Control Release* 2006; **116**: 1-27 [PMID: 17050027 DOI: 10.1016/j.jconrel.2006.08.013]
- 57 **Chaudhury A**, Das S. Recent advancement of chitosan-based nanoparticles for oral controlled delivery of insulin and other therapeutic agents. *AAPS PharmSciTech* 2011; **12**: 10-20 [PMID: 21153572 DOI: 10.1208/s12249-010-9561-2]
- 58 **Vllasaliu D**, Exposito-Harris R, Heras A, Casettari L, Garnett M, Illum L, Stolnik S. Tight junction modulation by chitosan nanoparticles: comparison with chitosan solution. *Int J Pharm* 2010; **400**: 183-193 [PMID: 20727955 DOI: 10.1016/j.ijpharm.2010.08.020]
- 59 **Sonaje K**, Lin KJ, Tseng MT, Wey SP, Su FY, Chuang EY, Hsu CW, Chen CT, Sung HW. Effects of chitosan-nanoparticle-mediated tight junction opening on the oral absorption of endotoxins. *Biomaterials* 2011; **32**: 8712-8721 [PMID: 21862121 DOI: 10.1016/j.biomaterials.2011.07.086]
- 60 **Bakhru SH**, Furtado S, Morello AP, Mathiowitz E. Oral delivery of proteins by biodegradable nanoparticles. *Adv Drug Deliv Rev* 2013; **65**: 811-821 [PMID: 23608641 DOI: 10.1016/j.addr.2013.04.006]
- 61 **Kadiyala I**, Loo Y, Roy K, Rice J, Leong KW. Transport of chitosan-DNA nanoparticles in human intestinal M-cell model versus normal intestinal enterocytes. *Eur J Pharm Sci* 2010; **39**: 103-109 [PMID: 19913612 DOI: 10.1016/j.ejps.2009.11.002]
- 62 **Douglas KL**, Piccirillo CA, Tabrizian M. Cell line-dependent internalization pathways and intracellular trafficking determine transfection efficiency of nanoparticle vectors. *Eur J Pharm Biopharm* 2008; **68**: 676-687 [PMID: 17945472 DOI: 10.1016/j.ejpb.2007.09.002]
- 63 **Richard I**, Thibault M, De Crescenzo G, Buschmann MD, Lavertu M. Ionization behavior of chitosan and chitosan-DNA polyplexes indicate that chitosan has a similar capability to induce a proton-sponge effect as PEI. *Biomacromolecules* 2013; **14**: 1732-1740 [PMID: 23675916 DOI: 10.1021/bm4000713]
- 64 **Guo S**, Huang L. Nanoparticles escaping RES and endosome: Challenges for siRNA delivery for cancer therapy. *J Nanomater* 2011; **2011**: 742895 [DOI: 10.1155/2011/742895]
- 65 **Köping-Höggård M**, Tubulekas I, Guan H, Edwards K, Nilsson M, Vårum KM, Artursson P. Chitosan as a nonviral gene delivery system. Structure-property relationships and characteristics compared with polyethylenimine in vitro and after lung administration in vivo. *Gene Ther* 2001; **8**: 1108-1121 [PMID: 11526458 DOI: 10.1038/sj.gt.3301492]
- 66 **Kim TH**, Ihm JE, Choi YJ, Nah JW, Cho CS. Efficient gene delivery by urocanic acid-modified chitosan. *J Control Release* 2003; **93**: 389-402 [PMID: 14644588 DOI: 10.1016/j.jconrel.2003.08.017]
- 67 **Chang KL**, Higuchi Y, Kawakami S, Yamashita F, Hashida M. Efficient gene transfection by histidine-modified chitosan through enhancement of endosomal escape. *Bioconjug Chem* 2010; **21**: 1087-1095 [PMID: 20499901 DOI: 10.1021/bc1000609]
- 68 **Kim TH**, Kim SI, Akaike T, Cho CS. Synergistic effect of poly(ethylenimine) on the transfection efficiency of galactosylated chitosan/DNA complexes. *J Control Release* 2005; **105**: 354-366 [PMID: 15949861 DOI: 10.1016/j.jconrel.2005.03.024]
- 69 **Parhamifar L**, Andersen H, Wu L, Hall A, Hudzech D, Moghimi SM. Polycation-mediated integrated cell death processes. *Adv Genet* 2014; **88**: 353-398 [PMID: 25409612 DOI: 10.1016/B978-0-12-800148-6.00012-2]
- 70 **Wong K**, Sun G, Zhang X, Dai H, Liu Y, He C, Leong KW. PEI-g-chitosan, a novel gene delivery system with transfection efficiency comparable to polyethylenimine in vitro and after liver administration in vivo. *Bioconjug Chem* 2006; **17**: 152-158 [PMID: 16417264 DOI: 10.1021/bc0501597]
- 71 **Jiang HL**, Kim YK, Arote R, Nah JW, Cho MH, Choi YJ, Akaike T, Cho CS. Chitosan-graft-polyethylenimine as a gene carrier. *J Control Release* 2007; **117**: 273-280 [PMID: 17166614 DOI: 10.1016/j.jconrel.2006.10.025]
- 72 **Fang N**, Chan V, Mao HQ, Leong KW. Interactions of phospholipid bilayer with chitosan: effect of molecular weight and pH.

- Biomacromolecules* 2001; **2**: 1161-1168 [PMID: 11777388 DOI: 10.1021/bm015548s]
- 73 **Huang M**, Fong CW, Khor E, Lim LY. Transfection efficiency of chitosan vectors: effect of polymer molecular weight and degree of deacetylation. *J Control Release* 2005; **106**: 391-406 [PMID: 15967533 DOI: 10.1016/j.jconrel.2005.05.004]
 - 74 **Mao HQ**, Roy K, Troung-Le VL, Janes KA, Lin KY, Wang Y, August JT, Leong KW. Chitosan-DNA nanoparticles as gene carriers: synthesis, characterization and transfection efficiency. *J Control Release* 2001; **70**: 399-421 [PMID: 11182210 DOI: 10.1016/S0168-3659(00)00361-8]
 - 75 **Kiang T**, Bright C, Cheung CY, Stayton PS, Hoffman AS, Leong KW. Formulation of chitosan-DNA nanoparticles with poly(propyl acrylic acid) enhances gene expression. *J Biomater Sci Polym Ed* 2004; **15**: 1405-1421 [PMID: 15648571 DOI: 10.1163/1568562042368112]
 - 76 **Wagner E**. Application of membrane-active peptides for nonviral gene delivery. *Adv Drug Deliv Rev* 1999; **38**: 279-289 [PMID: 10837761 DOI: 10.1016/S0169-409X(99)00033-2]
 - 77 **Li W**, Nicol F, Szoka FC. GALA: a designed synthetic pH-responsive amphipathic peptide with applications in drug and gene delivery. *Adv Drug Deliv Rev* 2004; **56**: 967-985 [PMID: 15066755 DOI: 10.1016/j.addr.2003.10.041]
 - 78 **Yang S**, May S. Release of cationic polymer-DNA complexes from the endosome: A theoretical investigation of the proton sponge hypothesis. *J Chem Phys* 2008; **129**: 185105 [PMID: 19045433 DOI: 10.1063/1.3009263]
 - 79 **Thibault M**, Astolfi M, Tran-Khanh N, Lavertu M, Darras V, Merzouki A, Buschmann MD. Excess polycation mediates efficient chitosan-based gene transfer by promoting lysosomal release of the polyplexes. *Biomaterials* 2011; **32**: 4639-4646 [PMID: 21450340 DOI: 10.1016/j.biomaterials.2011.03.010]
 - 80 **Thibault M**, Nimesh S, Lavertu M, Buschmann MD. Intracellular trafficking and decondensation kinetics of chitosan-pDNA polyplexes. *Mol Ther* 2010; **18**: 1787-1795 [PMID: 20628361 DOI: 10.1038/mt.2010.143]
 - 81 **Pichon C**, Billiet L, Midoux P. Chemical vectors for gene delivery: uptake and intracellular trafficking. *Curr Opin Biotechnol* 2010; **21**: 640-645 [PMID: 20674331 DOI: 10.1016/j.copbio.2010.07.003]
 - 82 **Lam AP**, Dean DA. Progress and prospects: nuclear import of nonviral vectors. *Gene Ther* 2010; **17**: 439-447 [PMID: 20200566 DOI: 10.1038/gt.2010.31]
 - 83 **Opanasopit P**, Rojanarata T, Apirakaramwong A, Ngawhirunpat T, Ruktanonchai U. Nuclear localization signal peptides enhance transfection efficiency of chitosan/DNA complexes. *Int J Pharm* 2009; **382**: 291-295 [PMID: 19716869 DOI: 10.1016/j.ijpharm.2009.08.029]
 - 84 **Zhao R**, Sun B, Liu T, Liu Y, Zhou S, Zuo A, Liang D. Optimize nuclear localization and intra-nucleus disassociation of the exogene for facilitating transfection efficacy of the chitosan. *Int J Pharm* 2011; **413**: 254-259 [PMID: 21536119 DOI: 10.1016/j.ijpharm.2011.04.039]
 - 85 **Brown BA**, Cooper EL. Immunological dichotomy in the larval bullfrog spleen. *Immunology* 1976; **30**: 299-305 [PMID: 770317]
 - 86 **Zaharoff DA**, Rogers CJ, Hance KW, Schlom J, Greiner JW. Chitosan solution enhances both humoral and cell-mediated immune responses to subcutaneous vaccination. *Vaccine* 2007; **25**: 2085-2094 [PMID: 17258843 DOI: 10.1016/j.vaccine.2006.11.034]
 - 87 **Yue Y**, Xu W, Xiong S. Modulation of immunogenicity and immunoprotection of mucosal vaccine against coxsackievirus B3 by optimizing the coadministration mode of lymphotactin adjuvant. *DNA Cell Biol* 2012; **31**: 479-488 [PMID: 21988406 DOI: 10.1089/dna.2011.1367]
 - 88 **Xia Y**, Fan Q, Hao D, Wu J, Ma G, Su Z. Chitosan-based mucosal adjuvants: Sunrise on the ocean. *Vaccine* 2015; **33**: 5997-6010 [PMID: 26271831 DOI: 10.1016/j.vaccine.2015.07.101]
 - 89 **van der Lubben IM**, Verhoef JC, van Aelst AC, Borchard G, Junginger HE. Chitosan microparticles for oral vaccination: preparation, characterization and preliminary in vivo uptake studies in murine Peyer's patches. *Biomaterials* 2001; **22**: 687-694 [PMID: 11246962 DOI: 10.1016/S0142-9612(00)00231-3]
 - 90 **Nagamoto T**, Hattori Y, Takayama K, Maitani Y. Novel chitosan particles and chitosan-coated emulsions inducing immune response via intranasal vaccine delivery. *Pharm Res* 2004; **21**: 671-674 [PMID: 15139524]
 - 91 **Goldmann K**, Ensminger SM, Spriewald BM. Oral gene application using chitosan-DNA nanoparticles induces transferable tolerance. *Clin Vaccine Immunol* 2012; **19**: 1758-1764 [PMID: 22933401 DOI: 10.1128/CI.00186-12]
 - 92 **Goldmann K**, Hoffmann J, Eckl S, Spriewald BM, Ensminger SM. Attenuation of transplant arteriosclerosis by oral feeding of major histocompatibility complex encoding chitosan-DNA nanoparticles. *Transpl Immunol* 2013; **28**: 9-13 [PMID: 23220147 DOI: 10.1016/j.trim.2012.11.007]
 - 93 **Wu ZH**, Ping QN, Lei XM, Li JY, Cai P. [Effects of the liposomes coated by chitosan and its derivatives on the gastrointestinal transit of insulin]. *Yaoxue Xuebao* 2005; **40**: 618-622 [PMID: 16196267]
 - 94 **Smoum R**, Rubinstein A, Srebnik M. Chitosan-pentaglycine-phenylboronic acid conjugate: a potential colon-specific platform for calcitonin. *Bioconjug Chem* 2006; **17**: 1000-1007 [PMID: 16848408 DOI: 10.1021/bc050357y]

P- Reviewer: Fernandes JC, Zhao ST **S- Editor:** Kong JX

L- Editor: A **E- Editor:** Lu YJ





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

